

# Steroid Production and Follicular Development of Neonatal Mouse Ovary during *In Vitro* Culture

Shabnam Abdi, M.Sc.<sup>1</sup>, Mojdeh Salehnia, Ph.D.<sup>1\*</sup>, Saman Hosseinkhani, Ph.D.<sup>2</sup>

1. Department of Anatomy, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

2. Department of Biochemistry, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran

## Abstract

**Background:** The aim of this study was to investigate follicular growth and steroid production in neonatal mouse ovary during *in vitro* culture.

**Materials and Methods:** In this experimental study, 7-day-old mouse whole ovaries were cultured in  $\alpha$ -MEM (medium supplemented with 100 mIU/ml recombinant follicle stimulating hormone, 1% insulin, transferrin and selenium (ITS), 5% fetal bovine serum (FBS), 100 IU/ml penicillin and 50  $\mu$ g/ml streptomycin for 7 days. The size of whole ovary was determined as mean area during culture. The survival rates of isolated preantral follicles after culture were assessed using trypan blue staining after being mechanically isolated. Histological evaluation of whole ovary was done by hematoxylin and eosin staining. 17- $\beta$  estradiol, progesterone and dehydroepiandrosterone concentrations in the medium were measured during culture.

**Results:** The mean area of ovary increased after culture (1.47 vs. 0.21 mm<sup>2</sup>). The survival rate of isolated follicles in ovary after culture was 99.2%. There was a significant decline in the percentage of primordial follicles after seven days of culture (91.8  $\pm$  0.2% vs. 65.1  $\pm$  1.1%), whereas the rate of preantral follicles increased significantly (4.6  $\pm$  0.4% vs. 29.2  $\pm$  0.5%). The levels of estradiol, progesterone and dehydroepiandrosterone also increased significantly after culture (p<0.001).

**Conclusion:** These results show that the growth and development of primordial follicles in contrast with hormonal production decreased during *in vitro* culture of neonatal mouse ovaries.

**Keywords:** *In vitro*, Primordial Follicles, Ovary, Organ Culture

**Citation:** Abdi Sh, Salehnia M, Hosseinkhani S. Steroid production and follicular development of neonatal mouse ovary during *In vitro* culture. *Int J Fertil Steril.* 2013; 7(3): 181-186.

## Introduction

Mammalian ovarian tissues contain a large number of resting pool of primordial follicles, which represent the reproductive potential of females. However, the number of follicles is limited and also gradually reduced during the female reproductive life (1, 2). Therefore, a considerably interesting issue in reproductive biology is how to initiate growth of these follicles to improve reproductive potential. *In vivo* and *in vitro* follicular growth and development are two alternative methods to initiate and activate their development within whole ova-

ries (3-6).

Almost 70 years ago techniques were used by Martinovitch (7) for *in vitro* culture of mammalian whole ovaries. Since then, *in vitro* organ or fragment culture has been evaluated under different culture conditions in several species, such as mice (6, 8-11), cattle (12, 13), baboon (14), bovine (15) and human (16, 17). Eppig and O'Brien in 1996 cultured newborn mouse ovaries for 8 days and activated primordial follicles and grew them to the secondary stage. The isolated secondary fol-

Received: 16 May 2012, Accepted: 17 Feb 2013

\* Corresponding Address: P.O. Box: 14115-111, Department of Anatomy, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Email: salehnm@modares.ac.ir



Royan Institute  
International Journal of Fertility and Sterility  
Vol 7, No 3, Oct-Dec 2013, Pages: 181-186

licles were cultured for 14 days, and mature and fertilizable oocytes and one living offspring were obtained (6).

O'Brien et al. (8) reported the development of oocytes derived from *in vitro* cultured primordial follicles isolated from cultured ovarian fragments and 59 living offsprings were produced.

The key advantage of this type of culture of ovarian fragments or whole ovary is that it preserves normal interactions between follicles at different sizes and stages and various extra follicular cell types within the ovarian tissue. However, the main disadvantage of this technique is that in large tissue fragments culture, degeneration and necrosis of tissues occur due to inadequate oxygenation. However additional investigations are required to improve this culture system.

The objective of the present study was to establish a mouse ovarian organ culture system where it could be used to evaluate a variety of factors which participate in folliculogenesis. During this organ culture, morphological and hormonal assessments of cultured ovaries were done.

## Materials and Methods

### *Animals and ovarian tissue*

In this experimental study, one week (n=15) and two weeks (n=5) old female mice obtained from National Medical Research Institute (NMRI) were housed and bred in the Central Animal House of Tarbiat Modares University. All animals were housed under a 12-hour light/12-hour dark regime at 22-24°C.

The mice were sacrificed by cervical dislocation and their ovaries were dissected free of fat and mesentery. For each one-week-old mouse, one ovary was selected randomly and fixed immediately in Bouin's solution (non-cultured control) and the other was considered for *in vitro* culture. Both ovaries of two-week-old mice were considered as control and fixed in Bouin's solution.

### *Organ culture*

Each ovary (n=15) was cultured in 24-well plates with tissue culture well inserts (non-tissue culture treated, PICM 012 50, 0.4- $\mu$ m pore size; Millipore Corp, Billerica, MA) in 0.4 ml  $\alpha$ -MEM

(Gibco, UK) supplemented with 5% fetal bovine serum (FBS), 100 mIU/ml recombinant follicle stimulating hormone (rFSH or Gonal-f, Serono, Switzerland), 1% insulin, transferrin and selenium (ITS, Gibco, UK), 100 IU/ml penicillin and 50  $\mu$ g/ml streptomycin. Approximately 400  $\mu$ l of culture medium was added to the compartment below the membrane insert, such that ovaries on the membrane were covered with a thin film of medium. The ovaries were incubated at 37°C and 5% CO<sub>2</sub> for 7 days. Every other day, 150  $\mu$ l media was replaced with fresh culture medium. The collected media were stored separately at -80°C until undertaking the hormonal assay.

### *Histological evaluation*

To assess the integrity of follicles after culturing, the follicular morphology was examined by histological staining. The non-cultured (control) ovaries from 7- and 14-day-old mice and cultured ovaries (for 7 days) from one-week-old mice were fixed in Bouin's solution and were embedded in paraffin wax and serially sectioned at 5- $\mu$ m-thickness (n=5 for each group); every five sections of each ovary were mounted on glass slides, and stained with hematoxylin and eosin. All sections were examined using light microscopy at a magnification of  $\times$ 400. For this study, stages of follicles have been classified and counted according to the method described previously.

### *Evaluation of the ovarian follicular viability using trypan blue*

The survival rates of the isolated preantral follicles from cultured ovaries (n=30 in each group) were determined using trypan blue staining. The preantral follicles with 120  $\mu$ m in diameter from the ovaries were mechanically isolated using insulin-gauge needles under stereomicroscope. Only follicles containing several layers of granulosa cells with a centrally located, healthy, visible oocyte and a thin layer of theca cells were selected. These isolated follicles were stained using trypan blue (0.4%) (Sigma, St. Louis, MO). The follicles were recorded as degenerated or survived: degenerated follicles stained blue and surviving ones not stained (13).

### *Ovarian area*

Photographs of each ovary in all groups of study

were prepared under Olympus (Tokyo, Japan) CK40 inverted microscope with an attached Olympus DP11 digital camera. All photographs were imported into ImageJ 1.33U software (National Institutes of Health, USA). Area was calculated in units of pixels and then converted to millimeter based on the pixel number to millimeter conversion ratio determined by measurements using the calibrated millimeter.

**Hormonal assays**

17-β estradiol (E2), progesterone (P4) and dehydroepiandrosterone (DHEA) were measured in collected media derived from ovarian culture during days 2, 4 and 6. The levels of 17-β estradiol (Monobind, USA, sensitivity=6.5 pg/mL) and progesterone (DiaPlus, USA, sensitivity=0.1 ng/ml) were measured by an enzyme immunoassay modified for the cell culture media. DHEA was measured with a commercial immunoenzymatic assay using antibodies directed against the α-subunit (Monobind, USA, sensitivity=20 pg/ml). These experiments were at minimum done in triplicates.

**Statistical analysis**


Statistical analysis was done with SPSS 16.0 software. The ovarian area and production of hormones were compared by one-way analysis of variance (ANOVA) and Tukey’s test. Student’s ttest was used to compare the proportion of follicular stages. P<0.05 was considered to be statistically significant.

**Results**


**Ovarian morphology**

The phase contrast morphology of ovaries is shown in fig 1. The appearance of growing follicles became apparent during the culture period and could be observed as swellings on the surface of the cultured ovary. The morphology of ovaries derived from 7-day-old mice before culture (non-cultured control) using the hematoxylin and eosin staining are shown in fig 2A, after one week culture in figs 2B and 2C and the ovaries of two-week-old mice in fig 2D. This figure showed that the structural organization of cultured ovaries was well preserved and appeared normal and it was similar to two-week-old mouse ovaries. Furthermore the necrosis area with atretic follicles (with a retracted oocyte and granulosa cells, pyknotic nu-

cleus) was consistently seen in the central part of the cultured ovary.



**Fig 1: Photomicrographs of 7-day-old mouse ovary viewed under the invert microscope in non-cultured fresh samples (A) and during one week of culture (B-D).**



**Fig 2: Hematoxylin and eosin staining of fresh and cultured mouse ovarian organ. Non-cultured fresh one week old mouse ovary which contains mainly primordial follicles with a few primary and secondary follicles (A), one week old mouse ovary after 7 days of culture (B). Higher magnification of one week old mouse ovary after 7 days, more secondary follicles were observed (C). Non-cultured fresh 14-day-old mouse ovary (D).**

**Normality rate of follicles**

The normality rates of follicles at various developmental stages within the ovaries of cultured and non-cultured controls are presented in fig 3. Ovaries derived from 7-day-old mice at the begin-

ning of the culture contained mostly primordial follicles ( $91.8 \pm 0.2\%$ ), with a small proportion of primary ( $5.8 \pm 2.5\%$ ) and secondary follicles ( $4.6 \pm 0.4\%$ ). After 7 days of *in vitro* organ culture, primordial follicles represented a smaller percentage of the total follicle pool ( $65.1 \pm 1.1\%$ ), similar to the follicle distribution seen in ovaries from 14-day-old mice ( $65.2 \pm 0.8\%$ ). The proportion of preantral follicles increased significantly during 7 days of culture, from  $4.6 \pm 0.4\%$  to  $29.2 \pm 0.5\%$  ( $p < 0.001$ ).




Fig 3: Normality rates of follicles after 7 days of *in vitro* ovarian culture. \*; significantly different with one week non-cultured ovary.

### The viability of isolated follicles

The morphologies of preantral follicles after trypan blue staining are shown in fig 4. The survival rates of preantral follicles derived from cultured ovaries using negative trypan blue staining was 99.2% and this rate was 100% in the control group of 14-day-old mice ovaries. There was no significant difference between them ( $p > 0.05$ ).




Fig 4: The trypan blue staining of preantral follicles. A. The survived follicle was not stained and B. degenerated follicle was stained intensively.

### Area of cultured ovaries

The area of cultured ovaries increased significantly from  $0.212 \pm 0.05 \text{ mm}^2$  on day 0 to  $1.47 \pm 0.1 \text{ mm}^2$  on day 7 of culture ( $p < 0.003$ , Fig 5).






Fig 5: Area of mouse ovary during 7 days of culture. \*; There were significant differences between lengths of culture.

### Hormonal assay

The levels of E2, P4 and DHEA during different lengths of culture are compared in fig 6. The levels of E2 in culture media on day 2, 4 and 6 were  $3381 \pm 43$ ,  $6552 \pm 214$  and  $12938 \pm 684 \text{ pg/ml}$  and the concentration of P4 on the same days were  $55.7 \pm 0.5$ ,  $111 \pm 3$  and  $157 \pm 14 \text{ ng/ml}$  respectively. DHEA levels increased from day 2 to day 4 to day 6 of culture ( $6.8 \pm 0.2$ ,  $14.5 \pm 0.2$  and  $29 \pm 0.5 \text{ ng/ml}$  respectively).





**Fig 6:** The level of estradiol (A), progesterone (B) and dehydroepiandrosterone (C) in collected culture media of mouse ovary. \*; There were significant differences between lengths of culture.

## Discussion

The establishment of a successful culture system for primordial follicle growth and development is essential to studies of *in vitro* follicle and ovarian culture and these kinds of studies are only at the incipient and nebulous stage.

Data obtained from this histological study showed that neonatal mouse ovary has a high percentage of primordial follicles. However, the proportion of these follicles decreased after 7 days of *in vitro* culture to a level observed in a 14-day-old mouse ovary. Concurrently, there was a sustained increase in the number of preantral follicles from day 2 to day 6 of culture.

Our observation confirmed that the ovarian culture conditions used in this study had a significant effect on follicular activation and growth. The culture of mouse whole ovary in  $\alpha$ -MEM medium supplemented with 10% FBS and 1% ITS not only increased the number of large follicles but also the ovarian area, implicating that an increase in ovarian area during culture may represent follicular growth and development.

In comparison with a previous work on the culture of isolated follicles (18), the whole ovary culture seems to yield better results. There are three possible explanations for the observed results.

First, *in vitro* ovarian culture provided follicles with an *in situ* growth environment resembling the ovary *in vivo* (19). The maintenance of the

follicle architecture supports the critical cellular interactions between adjacent somatic cells and between somatic and germ cells, which preserves local biochemical control pathways that trigger the initiation of follicular growth. In similar results, Jin SY et al. (20) demonstrated that culture of 8-day-old mouse ovaries for 4 days resulted in transition of the follicle population from primordial and primary to secondary follicles. These studies firstly establish that the activation of primordial follicles occur spontaneously, without the addition of growth factors or hormones. Second, the ovarian culture on tissue culture inserts covered by a thin film of medium survived better than isolated follicles. Third, enzymatic or mechanical dissociation of small follicles may have harmful effects on the subsequent follicular development (18).

Ovarian steroid production is an indicator of ovarian development and function. During the primordial follicle transition to the preantral follicle, granulosa and theca cells synthesize estradiol and androstenedione progressively, therefore measuring E2, P4 and DHEA could be an appropriate tool to evaluate the functionality of these cells within the ovary. The measurable steroidogenic function of these cells may be due to the remarkable increase in proliferation and differentiation of follicle cells and aromatase activities in granulosa cells during culture (21, 22).

## Conclusion

Results of this study show that *in vitro* ovarian organ culture could induce growth and development of follicles and steroid production, which indicate normal physiologic endocrine function of ovarian follicles during culture. Moreover ovarian culture alone is not able to complete the full growth and development of follicles to potentiate the competence of oocytes for fertilization. Thus, further research is needed to study *in vitro* culture of isolated follicles from ovarian culture. This study brings science one step closer to enhanced *in vitro* follicle culture methods, which may ultimately lead to clinical application in fertility treatments.

## Acknowledgements

This work was supported by Tarbiat Modares

University and Iran National Science Foundation.  
There is no conflict of interest in this study.

---

## References

1. Skinner, MK. Regulation of primordial follicle assembly and development. *Hum Reprod Update*. 2005; 11(5): 461-471.
  2. Hirshfield AN. Development of follicles in the mammalian ovary. *Int Rev Cytol*. 1991; 124: 43-101.
  3. Fortune, JE. The early stages of follicular development: activation of primordial follicles and growth of preantral follicles. *Anim Reprod Sci*. 2003; 78 (3-4): 135-163.
  4. Blandau RJ, Warrick E, Rumery RE. In vitro cultivation of fetal mouse ovaries. *Fertil Steril*. 1965; 16(6): 705-715.
  5. Blandau RJ, Odor DL. Observations on the behavior of oogonia and oocytes in tissue and organ culture. In: Biggers JD, Schuetz AW, editors. *Oogenesis*. Baltimore: University Park Press; 1972; 301-320.
  6. Eppig JJ, O'Brien ML. Development in vitro of mouse oocytes from primordial follicles. *Biol Reprod*. 1996; 54(1): 197-207.
  7. Martinovitch, PN. The development in vitro of the mammalian gonad. *Ovary and ovogenesis*. *Proc Roy Soc B*. 1938; 125: 232-249.
  8. O'Brien MJ, Pendola JK, Eppig JJ. A revised protocol for in vitro development of mouse oocytes from primordial follicles dramatically improves their developmental competence. *Biol Reprod*. 2003; 68(5): 1682-1686.
  9. Cortvrindt R, Smits J. Early preantral mouse follicle in vitro maturation: oocyte growth, meiotic maturation and granulosa-cell proliferation. *Theriogenology*. 1998; 49(4): 845-859.
  10. Klinger FG, De Felici M. In vitro development of growing oocytes from fetal mouse oocytes: stage-specific regulation by stem cell factor and granulosa cells. *Dev Biol*. 2002; 244(1): 85-95.
  11. Obata Y, Kono T, Hatada I. Gene silencing: maturation of mouse fetal germ cells in vitro. *Nature*. 2002; 418(6897): 497.
  12. Cecconi S, Barboni B, Coccia M, Mattioli M. In vitro development of sheep preantral follicles. *Biol Reprod*. 1999; 60(3): 594-601.
  13. Greenwald GS, Moor RM. Isolation and preliminary characterization of pig primordial follicles. *J Reprod Fertil*. 1989; 87(2): 561-571.
  14. Wandji SA, Srsen V, Nathanielsz PW, Eppig JJ, Fortune JE. Initiation of growth of baboon primordial follicles in vitro. *Hum Reprod*. 1997; 12(9): 1993-2001.
  15. Ralph JH, Wilmut I, Telfer EE. In vitro growth of bovine preantral follicles and the influence of FSH on follicle and oocyte diameter. *J Reprod Fertil Abstr Ser*. 1995; 15: 6.
  16. Hreinsson JG, Scott JE, Rasmussen C, Swahn ML, Hsueh AJ, Hovatta O. Growth differentiation factor-9 promotes the growth, development, and survival of human ovarian follicles in organ culture. *J Clin Endocrinol Metab*. 2002; 87(1): 316-321.
  17. Wright CS, Hovatta O, Margara R, Trew G, Winston RM, Franks S, et al. Effects of follicle-stimulating hormone and serum substitution on the in vitro growth of human ovarian follicles. *Hum Reprod*. 1999; 14(6): 1555-1562.
  18. Wandji SA, Eppig JJ, Fortune JE. FSH and growth factors affect the growth and endocrine function in vitro of granulosa cells of bovine preantral follicles. *Theriogenology*. 1996; 45(4): 817-832.
  19. Devine PJ, Rajapaksa KS, Hoyer PB. In vitro ovarian tissue and organ culture: a review. *Front Biosci*. 2002; 7: d1979-89.
  20. Jin SY, Lei L, Shikanov A, Shea LD, Woodruff TK. A novel two-step strategy for in vitro culture of early-stage ovarian follicles in the mouse. *Fertil Steril*. 2010; 93(8): 2633-2639.
  21. Peng X, Yang M, Wang L, Tong C, Guo Z. In vitro culture of sheep lamb ovarian cortical tissue in a sequential culture medium. *J Assist Reprod Genet*. 2010; 27(5): 247-257.
  22. Yang MY, Fortune JE. The capacity of primordial follicles in fetal bovine ovaries to initiate growth in vitro develops during mid-gestation and is associated with meiotic arrest of oocytes. *Biol Reprod*. 2008; 78(6): 1153-1161.
-