

# Reactive Oxygen Species and Antioxidant in Seminal Plasma and Their Impact on Male Fertility

Mohammad Eid Hammadeh, Ph.D.,<sup>1\*</sup>, Alixides Filippou A, M.Sc.<sup>1</sup>, Mohammed Faiz Hamad, M.Sc.<sup>1,2</sup>

1. Obstetrics and Gynaecology Department, IVF and Andrology Laboratory, University of Saarland, Humberg-Saar, Germany

2. Faculty of Pharmacy and Medical Technology, University of Petra, Amman, Jordan

## Abstract

Spermatozoa generate reactive oxygen species (ROS) in physiological amounts, which play a role in sperm functions during sperm capacitation, acrosome reaction (AR), and oocyte fusion. In addition, damaged sperm are likely to be the source of ROS. The most important ROS produced by human sperm are hydrogen peroxide, superoxide anion and hydroxyl radicals. Besides, human seminal plasma and sperm possess an antioxidant system to scavenge ROS and prevent ROS related cellular damage. Under normal circumstances, there is an appropriate balance between oxidants and antioxidants. A shift in the levels of ROS towards pro-oxidants in semen can induce oxidative stress (OS) on spermatozoa.

Male infertility is associated with increased ROS and decreased total antioxidant activity in the seminal plasma. ROS induce nuclear DNA strand breaks. Besides, due to a high polyunsaturated fatty acid content human sperm plasma membranes are highly sensitive to ROS induced lipid peroxidation thus decreasing membrane fluidity. This will result in increased lipid peroxidation (LPO), decreased sperm motility, viability, function and ultimately lead to infertility. The protective action of antioxidants against the deleterious effect of ROS on cellular lipids, proteins and DNA has been supported by several scientific studies.

The purpose of the present review is to address the possible relationship between ROS and antioxidants production in seminal plasma, and the role they may play in influencing the outcome of assisted reproductive technology (ART).

**Keywords:** Reactive Oxygen Species, Total Antioxidants, Infertility, Assisted Reproduction Technology, *In Vitro* Fertilization, Intracytoplasmic Sperm Injection

## Introduction

### *Infinity of infertility*

Since the first appearance of humans on earth, infertility has been one of the most controversial medical and social issues. Some civilizations considered it to be a punishment, while others thought of it as an illness. Some blamed it on the female; others could not explain it. This charge of infertility on the female, had serious consequences on her social image and psychological state (1). Additionally, physicians' lack of knowledge of gonadal and sperm function has been a main factor for the female to be considered the one responsible for infertility. It was not until the last decade that our knowledge of the human reproductive system allowed us to determine that a very important parameter of a couple's infertility has been male infertility and more specifically, sperm malfunction (2).

### *Normal and impaired sperm function*

Normal spermatozoa are those that successfully undergo a number of steps necessary for oocyte fertilization. The first step is maturation of the spermatozoa, which is initiated in the male genital tract and concludes with capacitation, the final step of maturation occurring in the female genital tract. Fully mature spermatozoa must swim in the female reproductive system, reach the oocyte, undergo acrosome reaction (AR), penetrate the zona pellucida and fuse with the oocyte pronucleus to form a zygote (3).

When, after successfully following all the above-mentioned steps, the spermatozoa are unable to lead to the natural fertilization of an oocyte, then impaired sperm function is considered. Failed attempts to conceive, after a period of two years, define infertility and the couple must visit an expert to seek diagnosis and possible treatment. The male

Received: 28 Jan 2009, Accepted: 27 Sep 2009

\* Corresponding Address: Obstetrics and Gynaecology Department, IVF and Andrology Laboratory, University of Saarland, 66421, Humberg-Saar, Germany  
Email: mehammadeh@yahoo.de



Royan Institute  
International Journal of Fertility and Sterility  
Vol 3, No 3, Nov-Dec 2009, Pages: 87-110

partner undergoes semen analysis and is evaluated according to the World Health Organization reference values. However, there are cases where the semen analysis indicates normal sperm, but the male is still infertile (4). This proves that there are parameters affecting sperm function that cannot be determined by routine semen analysis. Semen analysis includes assessment of seminal volume, spermatozoal motility, density, viability and morphology. Oxidative damage to spermatozoa, induced by excessive production of free radicals or impairment of the natural antioxidant mechanisms, has been identified as such a parameter (5).

### ***Free radicals: origin and oxidative stress Origin***

By definition, a free radical is any chemical compound with one or more unpaired electrons. The free radicals that have been associated with infertility are oxygen and oxygen-derived oxidants, namely, the superoxide anion ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ), peroxy radicals ( $ROO^\cdot$ ) and hydroxyl radicals ( $OH^\cdot$ ) (6, 7). These oxidants are widely known as reactive oxygen species (ROS) and, due to unpaired electron(s) tend to strongly react with other chemical compounds (8). More specifically, they seek stability by "stealing" electrons from nucleic acids, lipids, and proteins; leading to the damage of cells and disease phenomena (8, 9).

ROS are produced in the body by mitochondria, phagocytes, arachidonate pathways and other physiological processes in which they act as vital signalling molecules. They are products of natural oxygen metabolism and represent approximately 1 to 2% of metabolized oxygen (10). The balance between production and disposal of oxidant molecules is essential for tissue homeostasis. Increased rate of free radical production or decreased rate of removal leads to free radical accumulation and cellular damage (11). Additionally, their production is induced by external factors, such as cigarette smoke, ultraviolet light radiation, and others (7). ROS have been associated with the pathology of numerous diseases such as neurodegenerative diseases (12), vascular disease (13-15), cancer, diabetes, periodontal diseases (16-19) and of course, human infertility (20-28).

### ***ROS and oxidative stress (OS)***

The free radicals are a part of an organism's chemical reactions, and are necessary signalling molecules, as well as having a vital role for the maturation processes of several structures. Most free radicals in biology fit within the broader category of ROS, which include not only oxygen-centred radicals such as the superoxide anion radical ( $O_2^-$ ), hydroxyl radical ( $OH^\cdot$ ) or nitric oxide ( $NO^\cdot$ ), but

also some potentially dangerous non-radical derivatives of oxygen, such as hydrogen peroxide ( $H_2O_2$ ), peroxy nitrite anion ( $ONOO^-$ ), and hypochlorous acid ( $HOCL$ ) (28). The most common ROS that have a potential implication in reproductive biology include superoxide anion ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ), peroxy ( $ROO^\cdot$ ), and hydroxyl ( $OH^\cdot$ ) radicals.

The free radical nitric oxide ( $NO^\cdot$ ) and peroxy nitrite anion ( $ONOO^-$ ) also appear to play significant roles in reproduction and fertility. Öztezcan et al. (29) indicate that  $ONOO^-$  might cause sperm dysfunction through an increase in lipid peroxidation (LPO) and total sulphhydryl group depletion. The assumption that free radicals can influence male fertility has received substantial scientific support (30). Many reports have indicated that high levels of ROS are detected in the semen samples of 25% to 40% of infertile men (31). However, small controlled amounts of ROS are vital for spermatozoa to develop normally and be capable of fertilization structures (30, 32, 33).

In addition, de Lamirande et al. and Zini et al. reported that  $H_2O_2$  and  $O_2^-$  promote sperm capacitation and AR.  $H_2O_2$  also promotes hyperactivation and oocyte fusion (34, 35). Hydrogen peroxide ( $H_2O_2$ ) and superoxide anion are of great importance to spermatozoa. They are necessary for controlling the tyrosine phosphorylation events associated with sperm capacitation (32, 36).

### ***ROS production by spermatozoa***

ROS are also undoubtedly produced by spermatozoa (37-40), mainly through their mitochondrial system (41), as well as round cells during the spermatogenic process and epithelial cells. In the human ejaculate ROS are mainly produced by leukocytes with marginal amounts produced by spermatozoa (42). The production of ROS in ejaculated spermatozoa is initiated in immature germ cells (39) and continues in the epididymis when the surface of the spermatozoa is remodelled. When the mitochondrial capsule is assembled, chromatin undergoes condensation and motility is acquired for the capacitation of spermatozoa (43, 44). However,  $O_2^-$  production by spermatozoa has been questioned (45) on the basis that no free radical signal can be detected by electron paramagnetic resonance (EPR) spectroscopy. Electron leakage from complexes I and II of the mitochondrial transport chain has been proposed as a source of superoxide in male gametes (46). Numerous studies have shown that human sperm exhibit the capacity to generate ROS such as superoxide anion, hydrogen peroxide, and hydroxyl radicals (47, 48). The production of ROS by human sperm is due to a membrane-bound nicotinamide adenine dinucleotide (NADH) oxidase system (47).

***Spontaneous generation under aerobic conditions***

Spermatozoa generate small amount of  $O_2^-$  and NO, which are both tightly related to the cAMP pathway in the control of human sperm capacitation and protein tyrosine phosphorylation (36, 49). Beckman et al., suggested that these two radicals could combine to form peroxy nitrite ( $ONOO^-$ ) (50). Peroxy nitrite is not a free radical because the unpaired electrons of NO and  $O_2^-$  have combined to form a new N-O bond in peroxy nitrite, but it is a strong one or two electron oxidant and nitrating agent (51). Nevertheless, generating controlled low amounts of endogenous ROS by spermatozoa play a significant role in inducing sperm capacitation, AR, and acquisition of sperm-fertilizing ability (30, 52).

***Retained cytoplasm (RC)***

The production of ROS is higher in spermatozoa that are either damaged or retain abnormal cytoplasmic inclusions (39).

Radical production has also been detected in immature germ cells (44). According to Iwasaki and Gagnon (53), the capacity for ROS production is significantly enhanced in abnormal spermatozoa, and those sperm cells with retention of residual cytoplasm. Huszar and Vigue have found that morphological irregularities of sperm are significantly correlated with high creatine kinase (CK) activity (54). ROS production by spermatozoa has been associated with midpiece abnormalities, retained cytoplasm, cytoplasmic droplets and spermatozoa immaturity (48, 55-57).

The primary product of the immature spermatozoa system generating free radicals appears to be the superoxide anion ( $O_2^-$ ), which secondarily dismutase's to  $H_2O_2$  through the catalytic action of superoxide dismutase (SOD) (58,59). Moreover, the most prevalent ROS, hydrogen peroxide ( $H_2O_2$ ) is synthesized from  $O_2$  by mammalian spermatozoa (60) by a two-stage reduction of superoxide ( $O_2^*$ ) by  $H^+$  as the intermediate product (61). The retention of residual cytoplasm in the sperm midpiece after spermination has been associated with excessive production of ROS by spermatozoa (55). The ROS levels are expected to rise at a faster pace and in greater intensity in sperm samples in the presence of cytoplasmic residues. Morphometric analysis of the amount of residual cytoplasm present in the sperm midpiece has revealed significant correlations with the production of ROS (55).

The enzyme glucose-6-phosphate dehydrogenase (G6PD), which is superfluously exhibited in sperm residual cytoplasm, generates NADN, which in turn stimulates ROS formation (33, 43, 47, 54, 55).

Sperm with cytoplasmic droplets show a higher cellular content of cytoplasmic enzymes, including G6PD. This enzyme is responsible for the flux of

glucose through the hexose monophosphate shunt and the associated generation of NADPH. It is theorized that nicotinamide adenine dinucleotide phosphate (NADPH) generated via this system serves as the major source of electrons responsible for the production of  $O_2^-$  by human spermatozoa. Therefore, the retention of residual cytoplasm creates a situation in which sufficient substrate would be available to support excessive NADPH-dependent ROS generation (33, 62). Spermatozoa may generate ROS in two ways:

1. NADH-oxidase system at the level of sperm membrane level (37) and
2. NADH-dependent oxido-reductase (diphorase) at the level of the mitochondria (38).

Besides, the biochemical markers of the cytoplasmic space, such as creatine kinase, are positively correlated with the induction of peroxidative damage (55, 56). Huszar and Vigue (56) have found a positive relationship between CK activity and the rate of LPO, as measured by malondialdehyde (MDA) formation in sperm fractions. Hallak et al. have found an inverse relationship between CK levels and sperm morphological forms and suggested that CK levels can be used as a reliable marker for sperm quality and fertilizing potential in subfertile men (63).

Excessive ROS production by immature, morphologically abnormal spermatozoa with cytoplasmic residues such as those confronted in teratozoospermic semen specimens may induce oxidative damage of mature spermatozoa during sperm migration from the seminiferous tubules to the epididymis and may be an important cause of male infertility (64). Immature spermatozoa are a well-characterized source of ROS and a negative correlation between ROS production and semen quality has been documented (39). The excessive generation of ROS by abnormal spermatozoa and by contaminating leukocytes (leukospermia) has been identified as one of the few defined aetiologies for male infertility.

Many reports point out that those biochemical markers of the cytoplasmic space, such as creatine kinase, are positively correlated with the induction of peroxidative damage (55, 56).

Only one third of ROS produced by spermatozoa is released extracellularly (41). In the case of oligozoospermic males whose spermatozoa generate particularly high levels of ROS, the source of cytotoxic oxygen radicals is frequently intracellular (37, 55). However, the main ROS producing sources are immature spermatozoa, especially those with cytoplasmic droplets at the midpiece and leukocytes (65).

***ROS production by leukocytes***

ROS are produced by leukocytes, which are

present in the male reproductive system and in the ejaculate, as a result of their role in immunological defense against pathogenic germs (26, 66). Leukocytes are present throughout the male reproductive tract and are found in almost every human ejaculate (67). The majority of leukocytes in semen are granulocytes (50-60%), followed by macrophages (20% to 30%) and T lymphocytes (2% to 5%) (4). Leukocytes play an important role in immune surveillance (68, 69) and phagocyte clearance of abnormal sperm (66). However, white blood cells (WBC's) in human semen are also capable of ROS generation (70, 71). Genital tract inflammation and an increased number of leukocytes in the ejaculate have been repeatedly associated with male subfertility and infertility (41, 72-74). Leukocytes are a particularly important source of OS in the ejaculates of patients exhibiting leukocytospermia secondary to infection or as consequence of paraplegia (75). Activated leukocytes can produce 100-fold higher amounts of ROS than nonactivated leukocytes (41). Leukocytes may be activated in response to a variety of stimuli, including inflammation and infection (76).

Whittington and Ford demonstrated that infiltrating leukocytes are the predominant source of ROS production in unspecified sperm preparations (77). Sperm damage from ROS that is produced by leukocytes occurs if seminal leukocyte concentrations are abnormally high such as in leukocytospermia (78), or by removing seminal plasma during sperm preparation for assisted reproduction (72).

Leukocytospermia may induce an alteration in sperm structure by means of excessive ROS production by activated granulocytes. It has been shown that leukocytospermia and excessive ROS levels are associated with an increase in chromatin alterations and DNA damage in sperm, as defined by the sperm chromatin structure assay (79).

In patient samples that generated detectable ROS, the ability of the spermatozoa to retain motility for 24 hours after preparation on a 40/80% Percoll gradient was negatively correlated ( $r = -0.310$ ,  $p < 0.05$ ) with basal ROS production (77). Besides, ROS production was also related to the outcome of *in vitro* sperm mucus penetration tests. Unstimulated levels of ROS production showed a significant ( $p < 0.05$ ) negative correlation with the number of progressively motile spermatozoa present in the mucus after 15 ( $r = -0.379$ ) and 60 minutes ( $r = -0.362$ ) (77). Wang et al. demonstrated that mitochondrial function was inhibited in the spermatozoa of infertile men and significantly correlated with sperm concentration and the level of ROS production.

### ***ROS and male fertility: physiological association of ROS***

#### ***Association with subfertility***

Only in the case of excessive ROS production or malfunction of the native antioxidant production mechanisms, do the free oxidants cause problems by putting several tissues under OS (80-83). ROS have been widely associated with both the etiopathogenesis in female infertility (6, 21, 84), as well as with male subfertility and impaired sperm function, but their exact role is still not clear.

It has been more than a decade that the possibility of ROS negatively affecting male infertility has been examined by scientists (30, 82) and there has been an increasing generation of scientific data supporting that view (85). More than 60 years ago MacLeod and Ross et al. described the loss of sperm motility, when the latter is exposed to oxygen at 38°C (86) and 36 years later the first association of OS and impaired sperm function came from the University of Cambridge (87). Since then, numerous scientific publications with very interesting data have come to light, presenting the role of ROS in sperm physiology and morphology (55, 88), and subsequently, the reproductive outcome.

The plasma membrane of spermatozoa contains high amounts of polyunsaturated fatty acids (PUFA), making them highly susceptible to damage by OS (89-91). Furthermore, the concentration of scavenging enzymes, like SOD or glutathione peroxidase (92), in spermatozoa cytoplasm is very low, making the effect of OS more severe (52, 90, 93, 94). Scientific data has been presented to support that ROS negatively affect sperm function by contributing to the occurrence of LPO (47, 91, 95).

Increased production of ROS or impaired action of antioxidant mechanisms can lead to surplus of ROS and hence, OS. However, semen analysis of infertile men showed that increased levels of ROS are a result of over-production of ROS, rather than decreased enrichment of the seminal plasma with antioxidants (96). ROS produced by leukocytes in seminal plasma, sperm itself and other sources and the effect of ROS on male fertility depending on its concentration in seminal plasma despite of their sources of production whether it is pathological or non-pathological source.

The increased production of ROS from immature spermatozoa has been demonstrated in recent studies, indicating a negative correlation between the percentage of normal spermatozoa and levels of ROS production in semen. These results were obtained after calculating ROS production in an ejaculated sperm gradient of immature and mature spermatozoa (based on WHO parameters (4)), presenting high and low ROS production respectively,

and immature germ cells, which also presented with a low production of ROS (39, 97). This excessive production of ROS from immature sperm could cause DNA damage in mature sperm within the male reproductive tract and, hence, account partly for male infertility. It is well documented that there is a negative correlation between defective sperm chromatin structure (DNA-break) and fertility, in *in vitro* fertilization (IVF) cycles (98). However, this condition does not seem mandatory for successful fertilization; as demonstrated by intracytoplasmic sperm injection (ICSI), where normal fertilization and pregnancy rates can be achieved with cells that have not completed spermiogenesis, such as epididymal and testicular spermatozoa (99).

The applied assisted reproduction techniques (ART), as their title indicates, aim at achieving fertilization that is not able to occur naturally. However, research studies have shown that repeated cycles of centrifugation in the process of sperm preparation can induce the production of ROS by spermatozoa. This means that improving the motility of sperm through the routine preparation does not necessarily mean that occasional DNA damage does not occur at the same time, endangering the outcome of fertilization (100-103).

Hammadeh et al., showed that ROS and total antioxidants (TAS) concentration in seminal plasma did not differ significantly between the patients undergoing IVF or ICSI therapy, however, negative correlation was found between ROS concentration in seminal plasma and sperm vitality, membrane integrity, sperm density, chromatin condensation, and DNA single strand breaks in both IVF and ICSI groups (104).

#### **Sperm motility and hyperactivation**

ROS hydrogen peroxide, in particular, plays a positive physiological role in sperm hyperactivation and capacitation. Low concentration of a NO releasing compound, has been shown to be beneficial to the maintenance of post-thaw human sperm motility and viability (105).

Besides, ROS production has been related to the outcome of *in vitro* sperm mucus penetration tests. ROS showed a significant ( $p < 0.05$ ) negative correlation with the number of progressively motile spermatozoa present in the mucus after 15 ( $r = -0.379$ ) and 60 ( $r = -0.362$ ) minutes (77).

Moreover, a highly significant correlation was found between oxidation of sperm DNA and reduced motility [106]. Oxidative stress affects the mean semen parameters (count, motility, morphology) and therefore, asthenozoospermia is probably the best indicator for oxidative stress in a routine semen analysis.

#### **Sperm capacitation**

The molecular basis of sperm capacitation is still unclear. Calcium uptake, an increase in cAMP concentration, a rise in intracellular pH, an efflux of cholesterol from sperm plasma membrane (107-108) and tyrosine phosphorylation of specific proteins have been shown to occur during this process (109-110).

Capacitation of spermatozoa may thus occur by different ROS, but it takes place specifically by  $H_2O_2$  following an increase in cAMP, activation of protein kinase A, and downstream tyrosine kinase activation (36).

It has also been demonstrated by many authors (52, 111, 112) that ROS, such as superoxide anion ( $O_2^-$ ), hydrogen peroxidase ( $H_2O_2$ ), and nitric oxide  $NO$  can induce sperm capacitation *in vitro*. Production of nitric oxide ( $NO$ ) by spermatozoa has also been reported (51) and may serve as an additional oxidant source.  $NO$  has also been suggested to be involved in the capacitation of spermatozoa but only in the presence of  $H_2O_2$  (113).

#### **AR**

ROS other than  $H_2O_2$ , such as nitric oxide and superoxide anion  $O_2^-$ , have been shown to promote sperm capacitation and AR (35, 95).

Stimulation of endogenous NADPH-dependent ROS generation in human sperm appears to regulate AR via tyrosine phosphorylation (114).

#### **Sperm egg binding**

Aitken et al. demonstrated that low levels of ROS enhance the ability of human spermatozoa to bind to the zona pellucida, an effect that was reversed by the addition of vitamin E. (47).

Treatment of human spermatozoa with low concentrations of  $NO$  releasing compounds like sodium nitroprusside (SNP;  $100^{-7}$  -  $100^{-8}$  M) in the capacitating medium increased the number of spermatozoa bound to the hemizona (115).

#### **Sperm egg fusion**

The production of low concentrations of hydrogen peroxidase ( $H_2O_2$  and  $O_2^-$ ) by spermatozoa may have a functional role in the signalling events controlling capacitation and sperm-oocytes fusion (34, 36, 109, 116).

However, poor sperm-oocytes fusion, *in vitro* fertilization experiment (48, 109) and standard IVF (117-120) are related to high ROS production.

Aitken suggests that high failure rate of sperm-oocyte fusion bioassay can be related to increased generation of lipoperoxidase. In a subset of infertile patients sperm were refractory to the second messenger signal generated by  $Ca$  ionophore, excessively generated ROS and also exhibited a

high failure rate in sperm-oocytes fusion bioassays (70).

### **ROS level and fertilization**

The redox status of human spermatozoa is likely to affect phosphorylation and adenosine triphosphate (ATP) generation with a profound influence on its fertilizing potential. However, one of the main features of sperm that might profoundly affect fertilization and subsequent development is chromatin structure (121). ROS generation during OS associated with the appearance of damage to the DNA, especially in infertile patients, leads to a high incidence of DNA strand breaks (80, 122). High levels of seminal ROS also impaired the sperm fertilizing capacity by DNA damage and apoptosis (21, 80).

Under normal circumstances, spermatozoa with damaged DNA would not participate in the fertilization process because of collateral peroxidative damage to the sperm plasma membrane (123).

### **ROS level and cleavage (embryo) quality**

The production of ROS may also be involved in bovine embryo development (124). Zorn et al. found that high seminal plasma ROS levels are associated with impaired sperm fertilizing ability and lower pregnancy rates after IVF. In ICSI, a negative association of ROS with embryo development to the blastocyst stage has been observed (125).


### **ROS level and pregnancy**

After IVF, fertilization and pregnancy rates were negatively associated with ROS levels. In ICSI, a negative association of ROS with embryo development to the blastocyst stage has been observed, and significant fewer ICSI derived embryos reached the morula-blastocyst stage on day four (125). In the IVF group (n=26), eleven pregnancies were achieved (42.3%) with a 47.8% implantation rate per embryo transferred. In ICSI patients (n=22), four pregnancies were achieved (18.2%) with a 40.7% implantation rate per embryo transferred as previously described (126).

### **ROS and sperm DNA damage: pathology of ROS**

Several studies have shown that OS caused by ROS production induces damage to the sperm's DNA, even though sperm DNA appears to be more resistant than other cell types such as somatic cells (127). Specifically, sperm DNA is protected by its advanced packaging on the one hand (128) and seminal plasma antioxidants on the other (129). OS-induced DNA fragmentation, which is widely observed in the spermatozoa of infertile men (130, 131), may not directly affect the fertilizing ability of spermatozoa but it directly affects their contri-

bution to normal embryonic development (132) and the mutational load of the embryo (32) (Fig 1). Evidence also supports that OS causes significant damage not only to the nuclear, but also to the mitochondrial DNA of human spermatozoa (133).



**Fig 1: Schematic representation of the possible relationship between OS in the male germ line and abnormalities in the development of the embryo and the health and well-being of the offspring (134).**

Along their way from the testis to the oocyte, spermatozoa encounter many factors and conditions that can target their DNA and potentially damage it. Defective packaging of chromatin in the nucleus, apoptosis and ROS are the causes that attract the most scientific interest (135, 136). Sperm DNA damage may also account for the loss of the ability to conceive naturally (137, 138) and for a number of cases of unexplained pregnancy loss (139). As far as ARTs are concerned, their association with DNA damage has been widely discussed and examined in a variety of studies, and extensively reviewed by Agarwal and Allamaneni (140).

### **OS-induced apoptosis**

The programmed death of eukaryotic cells, which occurs without inducing an inflammatory response, is called apoptosis (141). Apoptosis during spermatogenesis has been assessed, discussed and supported in several studies (142-144) and has been associated with male infertility (142, 145, 146). However, research on apoptosis in ejaculated spermatozoa seems to still be in its developing stages, only discussed by a few studies (131, 147, 148), the latest primary article being that of Wang et al. (149). More specifically, the latter group reported a positive correlation between the presence of apoptotic markers, due to OS-induced apoptosis, and spermatozoa DNA damage. OS-induced apoptosis in ejaculated spermatozoa has also been reviewed recently by Agarwal and Said (150).

### **Role of NADPH and NADPH oxidase activity**

ROS are the intermediate steps of oxygen reduc-

tion, namely,  $O_2^-$ ,  $H_2O_2$  and  $OH$  are the products of reduction by one, two or three electrons, respectively. Also,  $HO_2$ , the acid of superoxide anion, is another free radical with a major contribution in the destructive LPO process occurring in spermatozoa (89). Exogenous molecules, such as aromatic derivatives and iron complexes with low molecular weight, can activate molecular oxygen by catalyzing electron transfer to it. This can also be done by activating  $\beta$ -nicotinamide adenine dinucleotide phosphate (NADPH) oxidase, which theoretically exists in human spermatozoa and is believed to transfer electrons from NADPH to ground state oxygen, leading to the formation of the superoxide anion radical. The latter is then dismutated to hydrogen peroxide ( $H_2O_2$ ), which is controlled by the antioxidant glutathione peroxidase. In case the function of the antioxidant is impaired for any reason, the spermatozoa experience hyperoxidation (OS status) (Fig 2) (24). Several other studies have supported the presence of NADPH oxidase-like activity, at the sperm plasma membrane, in human spermatozoa and the role of NADPH in  $O_2^-$  production and the subsequent consequences of induced OS on spermatozoa (33, 44, 46, 151, 152). The second ROS-generating system that has been proposed is a sperm diaphorase system, located in the middle piece and integrated into the mitochondrial respiratory system of the spermatozoa (33).

**Leukocytospermia and male infertility**

Apart from their role as members of the immune-defensive mechanism, leukocytes are also vital for the clearance of defective sperm via phagocytosis (66). Leukocytospermia is a condition which, according to WHO guidelines (4), is determined by the existence of a concentration greater than  $1 \times 10^6/ml$  of peroxidase-positive leukocytes in semen and it is encountered in an average of 15% of infertile men (73, 79, 94). Nevertheless, whether or not it is a pathological condition leading to abnormal morphology and impaired sperm function is controversial among scientists. Specifically, there are studies that have found a negative correlation and other studies that have

found a positive correlation of poor sperm quality/function, with increased WBC concentration (leukocytospermia) (74). More specifically, Curi et al. concluded that there is no positive correlation between leukocytospermia and impaired sperm motility (asthenozoospermia) (4, 153). In another study, Tomlinson et al. could not find any association of leukocyte concentration, neither with impaired sperm quality, nor with conception rates (154). Also, no correlation was found between leukocyte counts, sperm density and motility, sperm antibodies and growth of micro-organisms by el-Demiry et al. (67).

On the other hand, Yanushpolsky et al. presented a positive association of increased seminal granulocyte concentrations with abnormal semen parameters of statistical and clinical significance (155). In addition, Arata de Bella-barba et al. showed that increased WBC were a usual phenomenon in the semen of infertile men and associated with semen of poor quality parameters (156). Moreover, Lemkecher et al. supported the positive correlation, but also associated leukocytospermia with increased DNA fragmentation (157). Aitken et al. also presented a positive correlation between leukocytospermia and impaired sperm function, when leukocytes were introduced into the sperm mixture after preparation. The same correlation, when assessed during IVF, was found to be negative (48). In contrast, Vicino et al. showed that increases in leukocyte concentration in semen can affect the results of both IVF and ICSI procedures (158).

Apart from sperm quality, the effect of leukocytospermia on sperm morphology is also a highly controversial issue between researchers. Hence similarly, several studies have presented a positive correlation between increased leukocyte concentration in semen and abnormal sperm morphology. Specifically, Berger et al. showed a positive correlation between morphologically normal sperm and sperm penetration ability, in contrast to morphologically abnormal sperm due to increased leukocyte concentration (159).




Fig 2: Schematic representation of NADPH oxidase activity (24)

Eggert-Kruse et al. presented a positive correlation between high rates of leukocytes of the round cells with increased morphological abnormality (160) and Yanushpolsky et al. presented statistically significant differences in sperm morphology and leukocyte concentration of double threshold according to the WHO criteria (155). Moreover, Thomas et al. and Menkveld and Kruger presented a positive correlation between WBCs in general and polymorphonuclear granulocytes (PMN) respectively, with morphological defects (161, 162). Finally, two years ago, a primary article by Aziz et al. presented a highly significant negative correlation between leukocytospermia and impaired sperm structural integrity (163).

On the contrary, two more studies presented a statistically insignificant association between increased concentrations of leukocytes and morphologically abnormal sperm (164, 165). To enhance the controversy concerning the role of leukocytes in abnormal sperm morphology, Kiessling et al. and Tomlinson et al. presented a positive correlation between increased leukocyte numbers and morphologically normal sperm, due to leukocytes' physiological function of abnormal-sperm phagocytosis (66, 68).

Lackner et al. (166) suggested that anti-inflammatory medication of male patients with bacterial leukocytospermia improved sperm

count and reduced leukocyte concentrations. Furthermore, Oliva and Multigner (167), after using antihistamine-like drug treatment, reported a significant decrease in leukocytes in semen and a significant increase in sperm motility and morphology.

Furthermore, Gambera et al. (168) evaluated the effects of treatment with rofecoxib, a cyclooxygenase-2 inhibitor, on sperm quality and pregnancy rate after intrauterine insemination (IUI) or monitored intercourse. They have shown a significant reduction in leukocyte concentrations, and an improvement of sperm motility and morphology, particularly the presence and shape of the acrosomal complex and tail structure.

Evaluation of the effect of cigarette smoking on antioxidant levels and the presence of leukocytospermia in infertile men showed that lower levels of SOD and catalase were seen in infertile patients compared with fertile donors. SOD was significantly correlated with sperm concentration and negatively correlated with leukocytospermia. In addition, leukocytospermia was inversely correlated with sperm motility. SOD levels were negatively related to cigarette smoking. They concluded that cigarette smoking may impair sperm motility and decrease the antioxidant activity (negative correlation with SOD) in the seminal plasma (169).

*Table 1: Some natural antioxidants (Adapted from 186).*

Non-enzymatic antioxidant molecules		
Antioxidant Molecule	Subcellular location	
Ascorbate (vitamin C)	Plastid; apoplast; cytosol; vacuole	
$\beta$ -Carotene	Plastid	
Glutathione, reduced (GSH)	Plastid; mitochondrion; cytosol	
Polyamines (e.g., putrescine, spermine)	Nucleus; plastid; mitochondrion; cytosol	
$\alpha$ -Tocopherol (vitamin E)	Cell and plastid membranes	
Zeaxanthin	Chloroplast	
Antioxidant enzymes		
Enzyme	EC number	Subcellular location
Ascorbate peroxidase	1.11.1.11	Plastid stroma and membranes
Proxidases (non-specific)	1.11.1.7	Cytosol; cell wall-bound
Catalase	1.11.1.6	Glyoxysome; peroxisome; cytosol; mitochondria
Superoxide dismutase (SOD)	1.15.1.1	Cytosol (Cu/ZnSOD); plastid (Cu/ZnSOD; FeSOD); mitochondrion (MnSOD); peroxisome
Dehydroascorbate reductase	1.8.5.1	Cytosol; plastid
Glutathione reductase	1.6.4.2	Mitochondrion; cytosol; plastid
Monodehydroascorbate reductase	1.6.5.4	Plastid stroma
Glutathione S-transferases	2.5.1.18	Cytosol; microsomal



### ***Antioxidants and male infertility***

#### ***Types, role and origin***

ROS are necessary for physiological functioning of spermatozoa, but they need to be controlled and their concentrations maintained at a level that is not deleterious to the cells.

This function is carried out by antioxidants (170) (Table 1) present in seminal plasma (26, 171-175). The most common antioxidants, that protect spermatozoa from excess concentrations of ROS and OS-induced damage and altogether represent the total antioxidant capacity (TAC) of seminal plasma are SOD (58, 176), catalase (CAT) (177), the glutathione (GSH) peroxidase system selenium and selenoproteins such as the phospholipids hydroperoxide glutathione peroxidase (PHGPx) and the glutathione reductase system (178, 179), vitamins A, C (180) and E (181), glutathione (182), spermin, thiols, urate (183, 184), albumin, taurine and hypotaurine (171), L-carnitine and zinc. Additionally, the possible protective antioxidant role of vitamins K and D have been reported in literature (185). These antioxidants work together as a wide protective network and many of them become radicals themselves while scavenging oxidants. The rest of the antioxidants in the network ensure that they are regenerated back to their original structures. For example, vitamin C and glutathione regenerate vitamin E. However, this antioxidant protection is not always strong enough to protect from OS. When the volume and distribution of the spermatozoa's cytoplasm is abnormal, antioxidant enzymes are unable to be hosted properly and therefore their defensive chain-breaking function is impaired (42, 87, 90).

A number of reports have discussed the possible origin of antioxidants in semen, but our knowledge is still limited. Some studies supported the testicular origin of semen antioxidants, while others have presented evidence that the source of antioxidant activity is post-testicular. More specifically, in 1994, Bauche et al. presented evidence that SOD is of testicular origin, protecting it from deleterious ROS concentrations (187). In 1998, research by Yeung et al. confirmed that the source of antioxidants is post-testicular, most probably the prostate and seminal vesicles (188). Four years later, Zini et al. improved that report, showing that antioxidants are post-testicular products and their probable role is the protection of ejaculated spermatozoa from OS such as that which occurs in the female reproductive tract (175). However, during spermatogenesis and epididymal storage, sperm must rely on epididymal/testicular antioxidants and their own intrinsic antioxidant capacity for protection. Testicular biopsies from men with varicocele have shown an increase in oxidative DNA damage with-

in spermatogonia and spermatocytes (189).

#### ***Association with subfertility: a remedy?***

ROS increase or antioxidant deficit have been associated with a number of pathological conditions by several scientific studies. The protective action of antioxidants against the deleterious effect of ROS on cellular lipids, proteins and DNA has been supported by several scientific studies (173). In this review, we separated the publications into three groups; namely, studies discussing natural (enzymatic), synthetic (non-enzymatic) and both, respectively.

Antioxidants, in general, are free radical scavengers that suppress the formation of ROS and/or oppose their action. There are many antioxidants in seminal fluid which can be divided into two groups: enzymatic like such as SOD, catalase (96), and glutathione peroxidase, (178) and nonenzymatic antioxidants such as ascorbate, urate, alpha-tocopherol, pyruvate, glutathione, taurine, and hypotaurine (93). Antioxidants that are present in the seminal fluid compensate for the deficiency of cytoplasmic enzyme in the spermatozoa (190). Spermatozoa themselves possess high concentrations of thiol groups, as well as smaller amounts of ascorbic acid, alpha-tocopherol, uric acid and GSH (179, 191, 192).

ROS production and TAC can be used as a marker of OS in seminal fluid and has been correlated with male infertility. Infertile men with male factor or idiopathic diagnoses had significantly lower ROS-TAC scores than controls (85). Said et al. suggested that abnormal sperm morphology combined with elevated ROS production may serve as a useful indicator of potential damage to sperm DNA (193).

Lisak et al. examined the effect of oxidation of protein and lipids by a thermochemiluminescence (TCL) analyzer and showed that thermochemiluminescence indices in seminal plasma closely correlate with sperm characteristics among patients with sperm disturbances (194).

Morphologically abnormal and immature spermatozoa that retain cytoplasmic residues in the mid-piece can be separated by a double-density gradient procedure (39, 97, 195). Recently, a study conducted by Mendiola et al. (196) to compare dietary habits in normospermic and oligoasthenoteratospermic patients found that frequent intake of lipophilic foods like meat products or milk may negatively affect semen quality in humans, whereas some fruits or vegetables may maintain or improve semen quality. In another study of the same group, to compare the effect of specific nutrient intake, they found that a low intake of antioxidant nutrients was associated with a poor semen quality

in men that attended an infertility clinic (197).


**Enzymatic antioxidants**

Human sperm contains the enzyme system comprising glutathione peroxidase (GPX), glutathione reductase (GRD); and their substrates, glutathione (GSH) and glutathione disulfide (GSSG) (59, 178, 179). This system functions as a defence against LPO in human sperm by reducing formed lipid hydroperoxides, which are very reactive and damaging to the plasma membrane; and to hydroxylipids, which are essentially inert (198).

SOD, catalase, and glutathione peroxidase are antioxidants that convert superoxide (O<sub>2</sub><sup>-</sup>) and peroxide (H<sub>2</sub>O<sub>2</sub>) radicals to form O<sub>2</sub> and H<sub>2</sub>O.

**Superoxide dismutase (SOD)**

One family of antioxidants used by organisms as a defence mechanism against oxidative damage, is located in the family of SOD (Fig 3) (199-201). One of the members of this family, namely, extracellular superoxide dismutase (EC-SOD), is a vital part of the defense against ROS-mediated tissue damage, and is located in the extracellular matrix of tissues, hence, the name EC-SOD (202, 203). SOD acts to scavenge the superoxide anion, which is produced by a one-electron reduction of an oxygen molecule and initiates a radical chain reaction.



**Fig 3:** The enzymatic dismutation of superoxide anion by SOD, such as EC-SOD. In addition, the reaction of superoxide with nitric oxide to produce peroxynitrite anion is depicted along with the products of peroxynitrite decay (Adapted from 204).

In addition, the reaction of superoxide with nitric oxide to produce peroxynitrite anion is depicted along with the products of peroxynitrite decay (adapted from 204).

In 2000, Potts et al. reported that seminal plasma, whose role is to protect spermatozoa from DNA damage and LPO, contains antioxidants (205). Three years later, Lamond et al. presented evidence that adding SOD to IVF media would protect sperm chromatin from breakdown (206).

Earlier, in 1992, Nonogaki et al. had shown that adding SOD to culture medium protected sperm viability and the embryo development *in vivo* and *in vitro* (207).

The SOD antilipoperoxidative defence system in human sperm relies almost entirely on the activity of a single enzyme, the Cu/ZN isoform (208). SOD protects against spontaneous O<sub>2</sub> toxicity and lipid peroxidase (209). SOD and catalase also remove (O<sub>2</sub><sup>-</sup>) generated by NADH-oxidase in neutrophils and can play an important role in protecting spermatozoa during genitourinary inflammation (210).

The steady-state concentration of superoxide is under the control of extracellular SOD, the inhibition of which (by copper chelating) leads to the rapid disappearance of thiols (211). A pivotal role of SOD in protection of testicular cells against heat stress-induced apoptosis has been demonstrated *in vivo* and *in vitro* (212, 213).

Murawski et al.(214) showed a positive correlation between SOD activity in seminal plasma and semen quality parameters, sperm concentration and overall motility. SOD activity in oligoasthenozoospermic patients was significantly lower compared to the activity found in normospermic men. They came to the conclusion that decreased seminal plasma scavenger antioxidant capacity, particularly in the form of low SOD activity, can be responsible for male infertility.

**Catalase**

Catalase is a well known antioxidant enzyme whose localization is limited to peroxisome. Catalase activity has also been determined in human spermatozoa and the seminal plasma of fertile and infertile males (177). However, a difference in the seminal catalase activity of asthenozoospermic and oligo-asthenozoospermic patients with hyper-viscosity has been shown (215).

Tavilani et al. (216) found an inverse correlation between activities of CAT and SOD in seminal plasma with an MDA content of spermatozoa from normozoospermic samples. However, such correlations observed between activities of CAT and GPX of seminal plasma with MDA content of spermatozoa of asthenozoospermic patients. Furthermore, they found positive correlations between total activities of CAT, SOD and GPX with total content of MDA in seminal plasma from normozoospermic samples. In asthenozoospermic samples, there were no such significant correlations. They suspected that under pathological conditions (e.g. asthenozoospermia) the activity of seminal antioxidant enzymes can not protect spermatozoa and may cause an increase of lipid peroxidation from spermatozoa.

**Protein and SH group (SH-group)**

A sulfhydryl group (SH-group) plays an important role in sperm metabolism and antioxidative defence. Nakamura et al. demonstrated that seminal plasma SOD, catalase, glutathione peroxidase and sulfhydryl group levels are significantly lower in infertile patients than those in controls, suggesting their relationship to male infertility (217).

**Glutathione**

Reduced glutathione is also part of the antioxidant system in the seminal fluid. Glutathione is a tripeptidyl molecule and present in either the reduced glutathione (GSH) or oxidized form glutathione (GSSG) by forming a disulfide bond between two molecules.

Lenzi et al. found that glutathione injection (600 mg/day, i.m.) for two months significantly improved the morphology and motility patterns of spermatozoa. Even the *in vitro* use of glutathione treatment increased the forward motility and migration from the pellet of spermatozoa obtained from leukospermia samples (218). Parinaud et al. showed *in vitro* enhancement of sperm motility using a migration salt solution containing glucose and glutathione as an antioxidant (219).

A highly significant decrease of mean levels of antioxidants in seminal plasma (glutathione, ascorbic acid and total antioxidant status) were found in oligozoospermic and azoospermic cases compared to normozoospermic controls. Whereas, malondialdehyde level was significantly elevated in oligozoospermic and azoospermic men (220-222).

**L,N-acetyl-cysteine**

Glutathione ( $\gamma$ -glutamyl-cystenyl-glycine) is a natural, highly effective reducing agent. L, N-acetyl-cysteine is a potential regulator of germ-cell death and is a well established inhibitor of physiological cell death in several systems. N-acetyl-L-cysteine, a reducing substance, has been shown to improve sperm motility *in vitro* together with a decrease of ROS levels in infertile patients with a high seminal level of ROS. L, N-acetyl-cysteine (0.1, 1 and 5 mg/ml) has a dose dependent effect in reducing ROS levels; the reduction was greater in patients with high levels of ROS than in those with low levels (223). L, N-acetyl-cysteine and/or a mixture of essential fatty acids and natural vitamins A and E reduced levels of 8-hydroxy-2'-deoxyguanosine (8-OH-dG), are considered as a markers of OS-induced sperm DNA damage (140).

N, L acetyl-L-cysteine, when given in concentrations of 125, 100, 50 and 25 mmol/L, suppressed germ cell death in a dose-dependent manner, how-

ever, more research is needed to validate its efficacy *in vivo* (224).

**Glutathione peroxidase**

Glutathione peroxidase and catalase activities are of prostatic and multi-glandular origin, respectively (188).

Glutathione peroxidase, a selenium-containing antioxidant enzyme with glutathione as the electron donor, removes peroxy radical from various peroxides including  $H_2O_2$  to improve sperm motility. Minor alterations in sperm membranes in selected cases of dyspermia can be reversed by glutathione (GSH) therapy (225).

The presence of glutathione may prevent the accumulation of peroxynitrite ( $ONOO^-$ ) to toxic levels and may convert  $ONOO^-$  to secondary products with protective properties (226).

Low levels of NADH and glutathione, as a result of the increased activity of glutathione peroxidase to remove metabolites of membrane LPO will further affect cellular  $Ca^{+2}$  homeostasis (178).

The selenium-containing enzyme glutathione peroxidase destroys peroxides before they can damage the cell membrane and interacts synergistically with vitamin E (227).

Glutathione peroxidase has been measured in seminal plasma and correlated with male infertility (228). Yeung et al. studied the GPX, GSH reductase, and SOD activities of normozoospermic patients and patients undergoing *in vitro* fertilization treatment and suggested that the origin of the GPX, GSH, reductase, SOD is neither testicular nor epididymal (188).

**Glutathione reductase**

Glutathione peroxidase is one of the principal antioxidant defence enzymes in human spermatozoa, but it requires oxidized glutathione to be reduced by glutathione reductase using NADPH generated in the pentose phosphate pathway. Glutathione reductase regenerates reduced GSH from its oxidized form (GSSG). GSH has a likely role in sperm nucleus decondensation. Also, intracellular GSH concentrations found to be lower when sperm morphology is severely impaired in both fertile and infertile males (229).

Williams and Ford (230), showed that the pentose phosphate pathway in human spermatozoa can respond dynamically to oxidative stress and that inhibiting glutathione reductase impairs the ability of sperm to resist lipid peroxidation. They concluded that the glutathione peroxidase-glutathione reductase-pentose phosphate pathway system is functional and provides an effective antioxidant defence in normal human spermatozoa.

### ***Alpha-glutamyl transpeptidase***

Alpha-glutamyl transpeptidase is present in the midpiece and acrosomal region of spermatozoa and it may further regulate the GSH content of oocytes at the time of sperm penetration (225).

### ***Non-enzymatic antioxidants***

Besides the enzymatic antioxidant system, there are numerous non-enzymatic low molecular mass antioxidants which are believed to be even more important scavengers than high molecular mass compounds (96).

### ***Nitric oxide***

NO is synthesized from L-arginine by a family of enzymes known as the NOS. NO is a molecule of great biological significance and has long been considered to play an important role in sperm physiology (231).

NO can act as a free radical scavenger inactivating (59, 232) and even inhibiting production of superoxide anion ( $O_2^{\cdot-}$ ) (233) which cause LPO, a process which leads to the functional impairment of spermatozoa (234). NO is produced directly by spermatozoa and constitutive NOS is present in two isoforms similar to those present in both endothelial (eNOS) and in brain (bNOS) cells. Besides, spermatozoa from normozoospermic samples appear to have greater amounts of NOS and higher amounts of NO production than those from asthenozoospermic samples (235).

### ***Albumin***

Human serum albumin (HSA) present in culture media protects antioxidants. HSA has a cysteine residue in position 34 which is not involved in a disulfide bond and may exist in a different oxidative state: as a fully reduced sulfhydryl group, as a mixed disulfide with cysteine, glutathione or homocysteine, for example, or in a higher oxidative state like sulfonic acid. Serum albumin is discussed as a marker for systemic OS (236).

Albumin is reported to exhibit an excellent ability to sustain sperm motility (152).

### ***Zinc***

Zinc is an element whose importance in the biological systems is undisputed. Its importance can be understood by considering that when it is deficient, severe pathological consequences occur, such as acrodermatitis enteropathica, a rare autosomal-recessive inheritable disease (237). As far as reproduction is concerned, this element has been shown to be highly important for conception, successful implantation and pregnancy outcome (238, 239). Zinc is present in high concentrations in the seminal fluid and there is evidence that it may act

*in vivo* as a scavenger of excessive  $O_2^{\cdot-}$  production by defective spermatozoa and/or leukocytes in semen after ejaculation (240).

There is evidence that zinc plays a vital role in the physiology of spermatozoa and spermatogenesis. Specifically, Bedwal and Bahuguna reported that this element decreases testicular weight and causes shrinkage of seminiferous tubules (241). Its potential role in sperm production, viability and prevention of spermatozoa degradation and sperm membrane stabilization has also been supported (242). After ejaculation, the present abnormal spermatozoa are sources of oxidants, namely, superoxide anions, which bind with zinc and reduce its concentration in seminal plasma. Thus, zinc is considered to be a vital antioxidant, guarding normal spermatozoa against superoxide anion-induced OS (41, 225). Chia et al. supported the research(?) that zinc concentration in seminal plasma is significantly correlated with sperm density, motility and viability (243).

The biological function of zinc and the characteristic features of zinc deficiency have been reviewed (244, 245).

Zinc is vital for spermatogenesis and for the development of primary and secondary sexual characteristics (246). Experimental zinc deficiency in humans leads reversibly to reduced sperm count combined with reduced serum testosterone (247). There is a link between zinc deficiency and oligospermia (248).

The total zinc content of mammalian semen is high, 800-3000  $\mu\text{m/g}$  of dry weight, and it has been demonstrated that zinc deficiency induces atrophy of the seminiferous tubules and causes failure of spermatogenesis in rats (249). It has been suggested that the zinc ion exchange takes place between the epididymal epithelium and sperm cells as they pass along the epididymal duct and it might be significant for the maturation process of rat sperm cells during their passage through the epididymis (250, 251).

Ali et al. (252) showed a significant difference in serum and seminal zinc levels in normospermic, oligospermic ( $p < 0.05$ ) and azospermic ( $p < 0.005$ ) subjects. Levels of zinc in seminal plasma is correlated positively with sperm count and correlated negatively with sperm motility in both normospermic and oligospermic group, whereas levels of seminal plasma zinc showed negative correlations with semen volume, pH and leucocytes concentration in all three groups (normospermic, oligospermic and azospermic).

They concluded that zinc may contribute to fertility through its significant effects on various semen parameters and may help in the investigation and treatment of infertile males.

Others studies demonstrated that both folate and zinc have antioxidant properties that counteract ROS. Folate, zinc, ROS and thiols affect apoptosis, which is important for sperm release, regulation of follicle atresia, degeneration of the corpus luteum and endometrial shedding (253).

However, no consistent associations were found between antioxidant or zinc intakes and sperm aneuploidy (254).

### ***Vitamins C/E***

The most important antioxidants in seminal fluid seem to be vitamins C and E (180, 181). The concentration of vitamin C in seminal plasma is ten times greater than in blood plasma (364 vs. 40  $\mu\text{mol/L}$ ) (191). During the nonenzymatic recycling pathway for vitamin E regeneration that occurs in membranes, vitamin C reduces chromanoxyl radicals to recycle vitamin E and eventually being itself consumed by the process (255). Vitamin C intake may reduce DNA strand breaks (256) in human lymphocytes. Dietary supplementation of vitamin C protects human sperm from endogenous oxidative DNA damage (257). Hughes et al. found that *in vitro* treatment of sperm with antioxidants (300 and 600  $\mu\text{M}$  ascorbic acid; 3 and 60  $\mu\text{M}$  alpha-tocopherol, and 400  $\mu\text{M}$  urate) reduce the magnitude of DNA damage as measured by the comet assay (258, 259).

Vitamin E is a term that encompasses a group of potent, lipid-soluble, chain-breaking antioxidants. Structural analyses have revealed that molecules with vitamin E antioxidant activity include four tocopherols ( $\alpha\beta\gamma\delta$ ) and four tocotrienols ( $\alpha\beta\gamma\delta$ ) (260). Vitamin E is known to readily reduce alkyl peroxy radicals of unsaturated lipids (261). It is a major chain-breaking antioxidant in the sperm membranes and it appears to have a dose-dependent protective effect (262).

Vitamin E is proven to be effective in preventing LPO and other radical-driven oxidative events (263). Vitamin E functions as a chain-breaking antioxidant that prevents the propagation of the free radical reaction (264, 265). Vitamin E prevents the loss of spermatogenesis in males and failure to retain the zygote in female rats (266). Many investigators have found a reduction in the concentration of oxidized DNA (8-hydroxy-2 deoxyguanosine) in sperm after antioxidant supplementation (130, 267).

Vitamin E inhibits LPO in membranes by scavenging peroxy ( $\text{RO}\cdot$ ) and alkoxy ( $\text{ROO}\cdot$ ) radicals. However, the ability of alpha-tocopherol to maintain a steady-state rate of peroxy radical reduction in the plasma membrane depends on the recycling of  $\alpha$ -tocopherol by external reducing agents such as ascorbate or thiols (227).

Semen vitamin E levels were not increased after three months of daily treatment with 400 mg vitamin E plus 500 mg vitamin C, although there was a marked increase in serum concentration. Ascorbic acid (vitamin C) is an important water-soluble antioxidant that reduces sulfhydryls, scavenger's free radicals, and protects against endogenous oxidative DNA damage (257).

Small doses of vitamin C (200 mg), have been shown to increase the seminal level of ascorbate in smokers from 5.6 to 13.1 mg/dl, which is similar to the level achieved (16.1 mg/dl) after 1000 mg of vitamin C (268). In infertile patients with a high level of oxidative DNA damage in spermatozoa, even the combination of vitamin C and E with glutathione induced only a slight increase in sperm concentration (130). Concurrent administration of vitamin C (350 mg/d) and E (250 mg) *in vivo* were not able to prevent sperm DNA damage occurring after ejaculation (258). Vitamin C may become a pro-oxidant when free transition metals are present (92, 269).

Oral supplementation with vitamin C has been shown to improve sperm counts, motility, and morphology in infertile patients (270).

Seminal ascorbic acid level was significantly lower in patients with leukospermia than patients with normal semen parameters. Interestingly, a significantly greater percentage of men with abnormal DNA fragmentation index (DFI) were observed in the patients with low levels of seminal ascorbic acid compared to those with normal or high levels of ascorbic acid (59% vs. 33%,  $p < 0.05$ ). Therefore, they came to the conclusion that insufficient seminal ascorbic acid was frequently associated with sperm DNA damage (222).

Vitamins C and E both have antioxidant properties and have been shown to be implicated in metabolic pathways of free radical scavenging and OS protection (271, 272). Vitamin E specifically interacts with Se-containing glutathione peroxidase to prevent the oxidative breakdown of tissue membranes (273) and has been shown to improve sperm motility when supplemented orally (274), as well as the *in vitro* function of spermatozoa (275). In 1996, Geva et al. showed that treatment with vitamin E can protect from LPO and the consequent changes to the sperm (276). A number of clinical studies have concluded that supplementation of vitamins C and E in specific combinations, results in protecting OS-induced sperm DNA damage. More specifically, vitamins cause a decrease in 8-OH-dG levels, which is thought to be a marker of sperm DNA damage induced by OS (140). Moreover, evidence supports that adding vitamins C and E in the medium during sperm preparation reduces hydrogen peroxide and, hence, protects

against OS (19). Supplementation of ascorbic acid (vitamin C) during an earlier study resulted in significantly improved sperm quality (268).

Nouri et al. (277) studied the relationship between levels of LPO (malondialdehyde, MDA) and vitamins E and C concentrations in the sperm and seminal plasma of asthenoteratozoospermic (AOT) and normozoospermic men and their effect on semen parameters. They found that the level of vitamins E and C in seminal plasma of normozoospermic were significantly higher in asthenoteratozoospermic males, whereas MDA levels were significantly higher in AOT in comparison to normozoospermic males.

### **Urate**

Ascorbate, urate and protein sulfhydryls are the major antioxidants present in seminal plasma, while GSH is practically undetectable (188).

The proposed protective function of uric acid against free radicals in human blood (278) has not been sufficiently investigated in seminal plasma. Very little data could be found in the literature on its presence in seminal plasma (279) and on its antioxidative defence properties (280). Gavella et al. found changes in total seminal plasma antioxidative capacity chiefly due to ascorbate and urate (183). Orally supplemented ascorbic acid is rapidly distributed in all body tissue with the highest concentration in seminal vesicles (281).

### **Selenium**

Selenium, in the form of selenocysteine, functions as the catalytic centre in the active sites of at least nine human enzymes, including four glutathione peroxidase antioxidant enzymes (282-284). It is well known that selenium is involved in the male reproductive process. In mature rats the testicular content of selenium increases greatly at the beginning of spermatogenesis (285). Selenium is also required for normal testicular development and spermatogenesis in rats (286). Rats with low selenium levels produced sperm with impaired motility and characteristic mid-piece damage (287, 288) indicating that selenium is necessary for normal sperm development.

The specific role of selenium in spermatogenesis appears to be related to the phospholipid hydroperoxide glutathione peroxidase, which is expressed depending on the developmental state of spermatids and seems to be converted into a structural component in the mid-piece of mature spermatozoa (289). Selenium deficiency is associated with impaired sperm motility, structural alteration of the mid-piece, and loss of the flagellum (290). In mature spermatozoa, selenium is largely restricted to the mitochondrial capsule, a keratin-like matrix

that embeds the helix of mitochondria in the sperm mid-piece (291).

Phospholipids hydroperoxide glutathione peroxidase is expressed at higher levels in rat testes than in any other tissue (292). Deficiencies of selenium or glutathione can lead to instability of the mid-piece of spermatozoa, resulting in defective motility (293, 294).

Phospholipid hydroperoxide glutathione peroxidase (PHGPx) is a unique intracellular enzyme. Its uniqueness lies in the fact that it has the ability to reduce intracellular membrane phospholipid hydroperoxides, hence, its name (295). The highest activity of PHGPx has so far been measured in the testis, with a significant difference from that of other tissues, such as the brain and liver (296, 297). Due to the fact that sperm cells have high numbers of polyunsaturated fatty acids, they depend on PHGPx (together with other scavenging systems) to protect them from OS and LPO (298) which is the reason its concentration in spermatozoa is relatively high (295).

In 2005, Greco et al. presented an interesting study, supporting the antioxidant role of vitamins C and E. More specifically, they detected a reduction in ejaculated spermatozoa DNA fragmentation after treatment with a combination of the two vitamins. The interesting aspect of this study was that it was the first study to present the direct effect of antioxidant treatment on sperm DNA integrity *in vivo* (299). Glutathione peroxidase has been shown to maintain sperm motility through scavenging of peroxy radicals ( $\text{ROO}^-$ ) (300) and also protect sperm from LPO (301). Additionally, GSH has been shown to be a vital antioxidant mechanism maintaining normal sperm motility (62, 174) and the ability of sperm to undergo AR (62, 174). In an earlier study, Lenzi et al. presented that glutathione therapy for two months had a statistically significant positive effect on sperm motility and morphology (302).

### **Treatments with antioxidants**

According to a study carried out on patients with high levels of DNA damage or LPO and with a history of recurrent pregnancy loss, it was shown that increasing the intake of antioxidant could result in an improvement in pregnancy outcomes (303). After 90 days of treatment with the immune-modulating and anti-oxidants (beta-glucan, papaya, lactoferrin, and vitamins C and E) in patients with asthenoteratozoospermia associated with leukocytosis, Piomboni et al. (304) found a significant reduction in seminal fluid leukocyte concentrations ( $2.2 \pm 0.9$  vs.  $0.9 \pm 0.2$ ) and consequently an increase in the percentage of morphologically normal sperm ( $17.0 \pm 5.2\%$  vs.  $29.8 \pm 6.5\%$ ), to-

tal progressive motility ( $19.0 \pm 7.8$  vs.  $34.8 \pm 6.8$ ), and chromatin integrity. In a prospective study performed on men with persistent oligospermia (5–20 million/ml) a statistically significant ( $p = 0.009$ ) increase in sperm count after antioxidant therapy was recorded (305). In addition, univariate logistic regression analysis showed that men treated with antioxidant therapy presented with the probability of having a normal sperm count that was 20-fold ( $OR = 20.1$ ;  $CI\ 95\% = 1.05-43.2$ ;  $p = 0.014$ ) higher than a untreated men. The authors (305) are of the opinion that antioxidant therapy based on a combination of N-acetylcysteine (NAC) and micronutrient (vitamins- minerals) supplementation can be helpful in improving the sperm count at least in a subset of oligospermic males.

Moreover, studying the effect of zinc therapy in asthenozoospermic men showed that zinc therapy alone, in combination with vitamin E or with vitamin E and C was associated with an increase of sperm parameters and decrease of OS, sperm apoptosis and sperm DNA fragmentation index (DFI) (306).

Sperm cryopreservation extender supplemented with melatonin, pyruvate and taurine as antioxidants can improve semen motility after a freeze thawing procedure (307).

## Conclusion

ROS have been associated with the pathology of numerous diseases. Small controlled amounts of ROS are vital for spermatozoa to develop into normal spermatozoa capable of fertilization structures. Hydrogen peroxide,  $H_2O_2$ , and  $O_2^-$  promote sperm capacitation and AR. In addition, hydrogen peroxide ( $H_2O_2$ ) promotes hyperactivation and oocyte fusion. Besides, hydrogen peroxide and superoxide anion are of great importance to spermatozoa, and are necessary for controlling the tyrosine phosphorylation events associated with sperm capacitation. Additionally, ROS are produced by leukocytes present in the male reproductive system and in the ejaculate, as a result of their role in the immunological defense against pathogenic germs. Only in case of excessive production of ROS or malfunction of the native antioxidant-production mechanisms, do the free oxidants cause problems by putting tissues under OS.

The function of controlling excessive ROS production is carried out by antioxidants that are present in the seminal plasma. A number of reports have discussed the possible origin of antioxidants in semen but our knowledge on that is still limited. Some of the studies supported the testicular origin of semen antioxidants, while others have presented evidence that the source of antioxidant activity is post-testicular.

A review of the literature shows that the evidence about the scavenging role of antioxidants and sperm protection is controversial. Some researchers have presented evidence of a defensive role of antioxidants, others have found that both deficient and excessive concentrations of antioxidants are deleterious for the sperm and others have shown the effect of antioxidants on some sperm parameters but not all of them. The existing scientific data concerning the scavenging ability of organic and inorganic antioxidants is very interesting but their actual origin and their mechanism of action in the male reproductive system and on ejaculated spermatozoa warrant further studies.

## References

1. Morice P, Josset P, Chapron C, Dubuisson JB. History of infertility. *Hum Reprod Update*. 1995; 1: 497-504.
2. Hull MG, Glazener CM, Kelly NJ, Conway DI, Foster PA, Hinton RA, et al. Population study of causes, treatment, and outcome of infertility. *Br Med J (Clin Res Ed)*. 1985; 291: 1693-1697.
3. Larson KL, De Jonge CJ, Barnes AM, Jost LK, Evenson DP. Sperm chromatin structure assay parameters as predictors of failed pregnancy following assisted reproductive techniques. *Hum Reprod*. 2000; 15: 1717-1722.
4. World Health Organization. WHO laboratory manual for the examination of human semen and sperm-cervical mucus interaction. 4<sup>th</sup> Edition. Cambridge: Cambridge Univ Press; 1999.
5. Sikka SC, Rajasekaran M, Hellstrom WJ. Role of oxidative stress and antioxidants in male infertility. *J Androl*. 1995; 16: 464-468.
6. Agarwal A, Gupta S, Sharma RK. Role of oxidative stress in female reproduction. *Reprod Biol Endocrinol*. 2005; 3: 28.
7. Bagchi K, Puri S. Free Radicals and Antioxidants in Health and Disease. *East Mediterr Health J*. 1998; 4: 350-360.
8. Attaran M, Pasqualotto E, Falcone T, Goldberg JM, Miller KF, Agarwal A, et al. The effect of follicular fluid reactive oxygen species on the outcome of in vitro fertilization. *Int J Fertil Womens Med*. 2000; 45: 314-320.
9. Pierce JD, Cackler AB, Arnett MG. Why should you care about free radicals? *RN*. 2004; 67: 38-42; quiz 43.
10. Fulbert JC, Cals MJ. Free radicals in clinical biology. Origin, pathogenic effect and defense Mechanisms. *Pathol Biol (Paris)*. 1992; 40: 66-77.
11. Berry EM, Kohen R. Is the biological antioxidant system integrated and regulated? *Med Hypotheses*. 1999; 53: 397-401.
12. Gibson GE, Huang HM. Mitochondrial enzymes and endoplasmic reticulum calcium stores as targets of oxidative stress in neurodegenerative diseases. *J Bioenerg Biomembr*. 2004; 36: 335-340.
13. Evans MD, Dizdaroglu M, Cooke MS. Oxidative DNA damage and disease: induction, repair and significance. *Mutat Res*. 2004; 567: 1-61.
14. Madamanchi NR, Vendrov A, Runge MS. Oxidative stress and vascular disease. *Arterioscler Thromb Vasc*

Biol. 2005; 25: 29-38.

15. Halliwell B. Antioxidants in human health and disease. *Annu Rev Nutr.* 1996; 16: 33-50.
16. Lantos J, Roth E, Czopf L, Nemes J, Gal I. Monitoring of plasma total antioxidant status in different diseases. *Acta Chir Hung.* 1997; 36: 188-189.
17. Moore S, Calder KA, Miller NJ, Rice-Evans CA. Antioxidant activity of saliva and periodontal disease. *Free Radic Res.* 1994; 21: 417-425.
18. Pinzani P, Petruzzi E, Orlando C, Gallai R, Serio M, Pazzagli M. Serum antioxidant capacity in healthy and diabetic subjects as determined by enhanced chemiluminescence. *J Biolumin Chemilumin.* 1998; 13: 321-325.
19. Agarwal A, Allamaneni SS. Role of free radicals in female reproductive diseases and assisted reproduction. *Reprod Biomed Online.* 2004, 9: 338-347.
20. Agarwal A, Gupta S, Sharma R. Oxidative stress and its implications in female infertility - a clinician's perspective. *Reprod Biomed Online.* 2005; 11: 641-650.
21. Agarwal A, Saleh RA. Role of oxidants in male infertility: rationale, significance, and treatment. *Urol Clin North Am.* 2002; 29: 817-827.
22. Aitken RJ, Baker MA. Oxidative stress and male reproductive biology. *Reprod Fertil Dev.* 2004; 16: 581-588.
23. Baker MA, Aitken RJ. Reactive oxygen species in spermatozoa: methods for monitoring and significance for the origins of genetic disease and infertility. *Reprod Biol Endocrinol.* 2005; 3: 67.
24. de Lamirande E, Jiang H, Zini A, Kodama H, Gagnon C. Reactive oxygen species and sperm physiology. *Rev Reprod.* 1997; 2: 48-54.
25. Henkel R. The impact of oxidants on sperm function. *Andrologia.* 2005; 37: 205-206.
26. Kim JG, Parthasarathy S. Oxidation and the spermatozoa. *Semin Reprod Endocrinol.* 1998; 16: 235-239.
27. Sanocka D, Kurpisz M. Reactive oxygen species and sperm cells. *Reprod Biol Endocrinol.* 2004; 2: 12.
28. Halliwell B. Free radicals and vascular disease: how much do we know? *BMJ.* 1993; 307: 885-886.
29. Oztezcan S, Turkoglu UM, Kervancioglu E, Kocak T, Kocak-Toker N, Aykac-Toker G. In vitro effects of peroxynitrite on human spermatozoa. *Andrologia.* 1999; 31: 195-198.
30. Gagnon C, Iwasaki A, De Lamirande E, Kovalski N. Reactive oxygen species and human spermatozoa. *Ann N Y Acad Sci.* 1991; 637: 436-444.
31. Padron OF, Brackett NL, Sharma RK, Lynne CM, Thomas AJ, Jr., Agarwal A. Seminal reactive oxygen species and sperm motility and morphology in men with spinal cord injury. *Fertil Steril.* 1997; 67: 1115-1120.
32. Aitken RJ. The Amoroso Lecture. The human spermatozoon—a cell in crisis? *J Reprod Fertil.* 1999; 115: 1-7.
33. Aitken RJ, Fisher HM, Fulton N, Gomez E, Knox W, Lewis B, et al. Reactive oxygen species generation by human spermatozoa is induced by exogenous NADPH and inhibited by the flavoprotein inhibitors diphenylene iodonium and quinacrine. *Mol Reprod Dev.* 1997; 47: 468-482.
34. de Lamirande E, Gagnon C. Human Sperm Hyperactivation and Capacitation as parts of an Oxidative Process. *Free Radic Biol Med.* 1993; 14: 255-265.
35. Zini A, de Lamirande E, Gagnon C. Lower Levels of Nitric Oxide promote Human Sperm Capacitation in vitro. *J Androl.* 1996; 16: 424-431.
36. Aitken RJ, Harkiss D, Knox W, Paterson M, Irvine DS. A novel signal transduction cascade in capacitating human spermatozoa characterised by a redox-regulated, cAMP-mediated induction of tyrosine phosphorylation. *J Cell Sci.* 1998; 111(Pt 5): 645-656.
37. Aitken RJ, Buckingham DW, West KM. Reactive oxygen species and human spermatozoa: analysis of the cellular mechanisms involved in luminol- and lucigenin-dependent chemiluminescence. *J Cell Physiol.* 1992, 151: 466-477.
38. Gavella M, Lipovac V. Pentoxifylline-mediated reduction of superoxide anion production by human spermatozoa. *Andrologia.* 1992; 24: 37-39.
39. Gil-Guzman E, Ollero M, Lopez MC, Sharma RK, Alvarez JG, Thomas AJ, Jr., et al. Differential production of reactive oxygen species by subsets of human spermatozoa at different stages of maturation. *Hum Reprod.* 2001; 16: 1922-1930.
40. Hendin BN, Kolettis PN, Sharma RK, Thomas AJ Jr, Agarwal A. Varicocele is associated with elevated spermatozoal reactive oxygen species production and diminished seminal plasma antioxidant capacity. *J Urol.* 1999; 161: 1831-1834.
41. Plante M, de Lamirande E, Gagnon C. Reactive oxygen species released by activated neutrophils, but not by deficient spermatozoa, are sufficient to affect normal sperm motility. *Fertil Steril.* 1994; 62: 387-393.
42. Aitken RJ, Baker MA: Reactive oxygen species generation by human spermatozoa: a continuing enigma. *Int J Androl.* 2002; 25: 191-194.
43. Agnihotri S, Purohit SB, Laloraya M, Kumar GP. Regional heterogeneity in intracellular distribution of superoxide and hydrogen peroxide within the sperm and its relation to sperm development. *Arch Androl.* 1999, 43:113-121.
44. Fisher HM, Aitken RJ: Comparative analysis of the ability of precursor germ cells and epididymal spermatozoa to generate reactive oxygen metabolites. *J Exp Zool.* 1997; 277: 390-400.
45. Richer SC, Ford WC. A critical investigation of NADPH oxidase activity in human spermatozoa. *Mol Hum Reprod.* 2001; 7: 237-244.
46. Vernet P, Fulton N, Wallace C, Aitken RJ. Analysis of reactive oxygen species generating systems in rat epididymal spermatozoa. *Biol Reprod.* 2001; 65: 1102-1113.
47. Aitken RJ, Clarkson JS, Fishel S. Generation of reactive oxygen species, lipid peroxidation, and human sperm function. *Biol Reprod.* 1989; 41: 183-197.
48. Aitken RJ, West K, Buckingham D. Leukocytic infiltration into the human ejaculate and its association with semen quality, oxidative stress, and sperm function. *J Androl.* 1994; 15: 343-352.
49. Belen Herrero M, Chatterjee S, Lefievre L, de Lamirande E, Gagnon C. Nitric oxide interacts with the cAMP pathway to modulate capacitation of human spermatozoa. *Free Radic Biol Med.* 2000; 29: 522-536.
50. Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA. Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci USA.* 1990; 87: 1620-1624.



51. Herrero MB, de Lamirande E, Gagnon C. Tyrosine function of peroxyne, the reaction product of nitric oxide and superoxide. *Mol Hum Reprod.* 2001; 7: 913-921.
52. de Lamirande E, Gagnon C. Impact of reactive oxygen species on spermatozoa: a balancing act between-beneficial and detrimental effects. *Hum Reprod.* 1995; 10 Suppl 1: 15-21.
53. Iwasaki A, Gagnon C. Formation of reactive oxygen species in spermatozoa of infertile patients. *Fertil Steril.* 1992; 57: 409-416.
54. Huszar G, Vigue L: Incomplete development of human spermatozoa is associated with increased creatine phosphokinase concentration and abnormal head morphology. *Mol Reprod Dev.* 1993; 34: 292-298.
55. Gomez E, Buckingham DW, Brindle J, Lanzafame F, Irvine DS, Aitken RJ. Development of an image analysis system to monitor the retention of residual cytoplasm by human spermatozoa: correlation with biochemical markers of the cytoplasmic space, oxidative stress, and sperm function. *J Androl.* 1996; 17: 276-287.
56. Huszar G, Vigue L. Correlation between the rate of lipid peroxidation and cellular maturity as measured by creatine kinase activity in human spermatozoa. *J Androl.* 1994; 15: 71-77.
57. Rao B, Soufir JC, Martin M, David G. Lipid peroxidation in human spermatozoa as related to midpiece abnormalities and motility. *Gamete Res.* 1989; 24: 127-134.
58. Aitken RJ, Krausz C. Oxidative stress, DNA damage and the Y chromosome. *Reproduction.* 2001; 122: 497-506.
59. Alvarez JG, Touchstone JC, Blasco L, Storey BT. Spontaneous lipid peroxidation and production of hydrogen peroxide and superoxide in human spermatozoa. Superoxide dismutase as major enzyme protectant against oxygen toxicity. *J Androl.* 1987; 8: 338-348.
60. Totic J, Walton A. Formation of hydrogen peroxide by spermatozoa and its inhibitory effect of respiration. *Nature.* 1946; 158.
61. Holland MK, Alvarez JG, Storey BT. Production of superoxide and activity of superoxide dismutase in rabbit epididymal spermatozoa. *Biol Reprod.* 1982; 27: 1109-1118.
62. Baumber J, Ball BA, Gravance CG, Medina V, Davies-Morel MC. The effect of reactive oxygen species on equine sperm motility, viability, acrosomal integrity, mitochondrial membrane potential, and membrane lipid peroxidation. *J Androl.* 2000; 21: 895-902.
63. Hallak J, Sharma RK, Pasqualotto FF, Ranganathan P, Thomas AJ Jr., Agarwal A. Creatine kinase as an indicator of sperm quality and maturity in men with oligospermia. *Urology.* 2001; 58: 446-451.
64. Sikka SC. Role of oxidative stress and antioxidants in andrology and assisted reproductive technology. *J Androl.* 2004; 25: 5-18.
65. Garrido N, Meseguer M, Simon C, Pellicer A, Remohi J. Pro-oxidative and anti-oxidative imbalance in human semen and its relation with male fertility. *Asian J Androl.* 2004; 6: 59-65.
66. Tomlinson MJ, White A, Barratt CL, Bolton AE, Cooke ID. The removal of morphologically abnormal sperm forms by phagocytes: a positive role for seminal leukocytes? *Hum Reprod.* 1992; 7: 517-522.
67. el-Demiry MI, Young H, Elton RA, Hargreave TB, James K, Chisholm GD. Leucocytes in the ejaculate from fertile and infertile men. *Br J Urol.* 1986; 58: 715-720.
68. Kiessling AA, Lamparelli N, Yin HZ, Seibel MM, Eyre RC. Semen leukocytes: friends or foes? *Fertil Steril.* 1995; 64: 196-198.
69. Pudney J, Anderson D. Organization of immunocompetent cells and their function in the male reproductive tract. In: Griffin P, Johnson P (editors). *Local immunity in reproductive tract tissues.* Oxford: Oxford University Press; 1993.
70. Aitken RJ, West KM. Analysis of the relationship between reactive oxygen species production and leukocyte infiltration in fractions of human semen separated on Percoll gradients. *Int J Androl.* 1990; 13: 433-451.
71. Kessopoulou E, Tomlinson MJ, Barratt CL, Bolton AE, Cooke ID. Origin of reactive oxygen species in human semen: spermatozoa or leukocytes? *J Reprod Fertil.* 1992; 94: 463-470.
72. Ochsendorf FR. Infections in the male genital tract and reactive oxygen species. *Hum Reprod Update.* 1999; 5: 399-420.
73. Sharma RK, Pasqualotto AE, Nelson DR, Thomas AJ Jr, Agarwal A. Relationship between seminal white blood cell counts and oxidative stress in men treated at an infertility clinic. *J Androl.* 2001; 22: 575-583.
74. Wolff H. The biologic significance of white blood cells in semen. *Fertil Steril.* 1995; 63: 1143-1157.
75. Sharma RK, Agarwal A. Role of reactive oxygen species in male infertility. *Urology.* 1996; 48: 835-850.
76. Pasqualotto FF, Sharma RK, Nelson DR, Thomas AJ, Agarwal A. Relationship between oxidative stress, semen characteristics, and clinical diagnosis in men undergoing infertility investigation. *Fertil Steril.* 2000; 73: 459-464.
77. Whittington K, Ford WC. Relative contribution of leukocytes and of spermatozoa to reactive oxygen species production in human sperm suspensions. *Int J Androl.* 1999; 22: 229-235.
78. Shekarriz M, Thomas AJ Jr, Agarwal A. Effects of time and sperm concentration on reactive oxygen species formation in human semen. *Arch Androl.* 1995; 34: 69-75.
79. Alvarez JG, Sharma RK, Ollero M, Saleh RA, Lopez MC, Thomas AJ Jr, Evenson DP, Agarwal A. Increased DNA damage in sperm from leukocytospermic semen samples as determined by the sperm chromatin structure assay. *Fertil Steril.* 2002; 78: 319-329.
80. Agarwal A, Said TM. Role of sperm chromatin abnormalities and DNA damage in male infertility. *Hum Reprod Update.* 2003; 9: 331-345.
81. Aitken RJ. The Founders Lecture. Human spermatozoa; Fruits of Creation and Seeds of Doubt. *Reprod Fertil Dev.* 2005; 16: 655-664.
82. Aitken RJ, Clarkson JS. Cellular basis of defective sperm function and its association with the genesis of reactive oxygen species by human spermatozoa. *J Reprod Fertil.* 1987; 81: 459-469.
83. Balercia G, Armeni T, Mantero F, Principato G, Regoli F. Total oxyradical scavenging capacity toward different reactive oxygen species in seminal plasma and sperm cells. *Clin Chem Lab Med.* 2003; 41: 13-19.
84. Agarwal A, Gupta S. Role of reactive oxygen spe-

- cies in female reproduction. Part I. Oxidative stress: a general overview. *Women Health*. 2005.
85. Sharma RK, Pasqualotto FF, Nelson DR, Thomas AJ Jr, Agarwal A. The reactive oxygen species-total antioxidant capacity score is a new measure of oxidative stress to predict male infertility. *Hum Reprod*. 1999; 14: 2801-2807.
  86. MacLeod J. The role of oxygen in the metabolism and motility of human spermatozoa. *Am J Physiol*. 1943; 138: 512-518.
  87. Jones R, Mann T, Sherins R. Peroxidative breakdown of phospholipids in human spermatozoa, spermicidal properties of fatty acid peroxides, and protective action of seminal plasma. *Fertil Steril*. 1979; 31: 531-537.
  88. Aziz N, Saleh RA, Sharma RK, Lewis-Jones I, Esfandiari N, Thomas AJ Jr, et al. Novel association between sperm reactive oxygen species production, sperm morphological defects, and the sperm deformity index. *Fertil Steril*. 2004; 81: 349-354.
  89. Alvarez JG, Storey BT. Differential incorporation of fatty acids into and peroxidative loss of fatty acids from phospholipids of human spermatozoa. *Mol Reprod Dev*. 1995; 42: 334-346.
  90. Lewis SE, Boyle PM, McKinney KA, Young IS, Thompson W. Total antioxidant capacity of seminal plasma is different in fertile and infertile men. *Fertil Steril*. 1995; 64: 868-870.
  91. Zalata AA, Ahmed AH, Allamaneni SS, Comhaire FH, Agarwal A. Relationship between acrosin activity of human spermatozoa and oxidative stress. *Asian J Androl*. 2004; 6: 313-318.
  92. Guerin P, El Mouatassim S, Menezo Y. Oxidative stress and protection against reactive oxygen species in the pre-implantation embryo and its surroundings. *Hum Reprod Update*. 2001; 7: 175-189.
  93. Saleh RA, Agarwal A. Oxidative stress and male infertility: from research bench to clinical practice. *J Androl*. 2002; 23: 737-752.
  94. Saleh RA, Agarwal A, Kandirali E, Sharma RK, Thomas AJ, Nada EA, et al. Leukocytospermia is associated with increased reactive oxygen species production by human spermatozoa. *Fertil Steril*. 2002; 78: 1215-1224.
  95. Griveau JF, Dumont E, Renard P, Callegari JP, Le Lannou D. Reactive oxygen species, lipid peroxidation and enzymatic defence systems in human spermatozoa. *J Reprod Fertil*. 1995; 103: 17-26.
  96. Zini A, de Lamirande E, Gagnon C. Reactive oxygen species in semen of infertile patients: levels of superoxide dismutase- and catalase-like activities in seminal plasma and spermatozoa. *Int J Androl*. 1993; 16: 183-188.
  97. Ollero M, Gil-Guzman E, Lopez MC, Sharma RK, Agarwal A, Larson K, et al. Characterization of subsets of human spermatozoa at different stages of maturation: implications in the diagnosis and treatment of male infertility. *Hum Reprod*. 2001; 16: 1912-1921.
  98. Hammadeh ME, Stieber M, Haidl G, Schmidt W. Association between sperm chromatin condensation, morphology based on strict criteria, and fertilization, cleavage and pregnancy rates in an IVF program. *Andrologia*. 1998; 30: 29-35.
  99. Hammadeh ME, Al-Hasani S, Stieber M, Rosenbaum P, Kupker D, Diedrich K, et al. The effect of chromatin condensation (AnilineBlue staining) and morphology (strict criteria) of Human spermatozoa on fertilization, cleavage and pregnancy rates in an intracytoplasmic sperm injection programme. *Hum Reprod*. 1996; 11: 2468.
  100. Agarwal A, Ikemoto I, Loughlin KR. Levels of reactive oxygen species before and after sperm preparation: comparison of swim-up and L4 filtration. *Arch Androl*. 1994; 32: 169-174.
  101. Agarwal A, Ikemoto I, Loughlin KR. Effect of sperm washing on levels of reactive oxygen species in semen. *Arch Androl*. 1994; 33: 157-162.
  102. Lopes S, Jurisicova A, Sun JG, Casper RF. Reactive oxygen species: potential cause for DNA fragmentation in human spermatozoa. *Hum Reprod*. 1998; 13: 896-900.
  103. Zini A, Finelli A, Phang D, Jarvi K. Influence of semen processing technique on human sperm DNA integrity. *Urology*. 2000; 56: 1081-1084.
  104. Hammadeh ME, Al Hasani S, Rosenbaum P, Schmidt W, Fischer Hammadeh C. Reactive oxygen species, total antioxidant concentration of seminal plasma and their effect on sperm parameters and outcome of IVF/ICSI patients. *Arch Gynecol Obstet*. 2008; 277: 515-526.
  105. Hellstrom W, Neal D. Diagnosis and therapy of male genital tract infections. *Infertility and Reproductive Medicine clinics of North America*. 1992; 3(2): 399-411.
  106. Kao SH, Chao HT, Chen HW, Hwang TI, Liao TL, Wei YH. Increase of oxidative stress in human sperm with lower motility. *Fertil Steril*. 2008; 8(11): 2178-2184.
  107. Visconti PE, Kopf GS. Regulation of protein phosphorylation during sperm capacitation. *Biol Reprod*. 1998; 59: 1-6.
  108. Yanagimachi R. Mammalian fertilization. In: Knobil E, Neill J (editors). *The Physiology of Reproduction*. New York: Raven Press: 1994; 189-317.
  109. Aitken RJ. Free radicals, lipid peroxidation and sperm function. *Reprod Fertil Dev*. 1995; 7: 659-668.
  110. Leclerc P, de Lamirande E, Gagnon C. Cyclic adenosine 3',5'-monophosphate-dependent regulation of protein tyrosine phosphorylation in relation to human sperm capacitation and motility. *Biol Reprod*. 1996; 55: 684-692.
  111. Griveau JF, Le Lannou D: Effects of antioxidants on human sperm preparation techniques. *Int J Androl*. 1994; 17: 225-231.
  112. Herrero MB, de Lamirande E, Gagnon C. Nitric oxide regulates human sperm capacitation and protein-tyrosine phosphorylation in vitro. *Biol Reprod*. 1999; 61: 575-581.
  113. Zini A, De Lamirande E, Gagnon C. Low levels of nitric oxide promote human sperm capacitation in vitro. *J Androl*. 1995; 16: 424-431.
  114. Leclerc P, de Lamirande E, Gagnon C. Regulation of protein-tyrosine phosphorylation and human sperm capacitation by reactive oxygen derivatives. *Free Radic Biol Med*. 1997; 22: 643-656.
  115. Sengoku K, Tamate K, Yoshida T, Takaoka Y, Miyamoto T, Ishikawa M. Effects of low concentrations of nitric oxide on the zona pellucida binding ability of human spermatozoa. *Fertil Steril*. 1998; 69: 522-527.
  116. De Lamirande E, Gagnon C. A positive role for the superoxide anion in triggering hyperactivation and capacitation of human spermatozoa. *Int J Androl*. 1993;

16: 21-25.

117. De Geyter C, De Geyter M, Schneider HP, Nieschlag E. Subnormal sperm parameters in conventional semen analysis are associated with discrepancies between fertilization and pregnancy rates in in-vitro fertilization and embryo transfer. *Int J Androl.* 1992; 15: 485-497.
118. Krausz C, Mills C, Rogers S, Tan SL, Aitken RJ. Stimulation of oxidant generation by human sperm suspensions using phorbol esters and formyl peptides: relationships with motility and fertilization in vitro. *Fertil Steril.* 1994; 62: 599-605.
119. Sukcharoen N, Keith J, Irvine DS, Aitken RJ. Predicting the fertilizing potential of human sperm suspensions in vitro: importance of sperm morphology and leukocyte contamination. *Fertil Steril.* 1995; 63: 1293-1300.
120. Sukcharoen N, Keith J, Irvine DS, Aitken RJ. Prediction of the in-vitro fertilization (IVF) potential of human spermatozoa using sperm function tests: the effect of the delay between testing and IVF. *Hum Reprod.* 1996; 11: 1030-1034.
121. Sakkas D, Urner F, Bizzaro D, Manicardi G, Bianchi PG, Shoukir Y, et al. Sperm nuclear DNA damage and altered chromatin structure: effect on fertilization and embryo development. *Hum Reprod.* 1998; 13 Suppl 4: 11-19.
122. Evenson DP, Larson KL, Jost LK. Sperm chromatin structure assay: its clinical use for detecting sperm DNA fragmentation in male infertility and comparisons with other techniques. *J Androl.* 2002; 23: 25-43.
123. Kodama H, Kuribayashi Y, Gagnon C. Effect of sperm lipid peroxidation on fertilization. *J Androl.* 1996; 17: 151-157.
124. Blondin P, Coenen K, Sirard MA. The impact of reactive oxygen species on bovine sperm fertilizing ability and oocyte maturation. *J Androl.* 1997; 18: 454-460.
125. Zorn B, Vidmar G, Meden-Vrtovec H. Seminal reactive oxygen species as predictors of fertilization, embryo quality and pregnancy rates after conventional in vitro fertilization and intracytoplasmic sperm injection. *Int J Androl.* 2003; 26: 279-285.
126. Hammadeh ME, Radwan M, Al-Hasani S, Micu R, Rosenbaum P, Lorenz M, et al. Comparison of reactive oxygen species concentration in seminal plasma and semen parameters in partners of pregnant and non-pregnant patients after IVF/ICSI. *Reprod BioMed Online.* 2006; 13(5): 696-706.
127. McKelvey-Martin VJ, Melia N, Walsh IK, Johnston SR, Hughes CM, Lewis SE, et al. Two potential clinical applications of the alkaline single-cell gel electrophoresis assay: (1). Human bladder washings and transitional cell carcinoma of the bladder; and (2). Human sperm and male infertility. *Mutat Res.* 1997; 375: 93-104.
128. Fuentes-Mascorro G, Serrano H, Rosado A. Sperm chromatin. *Arch Androl.* 2000; 45: 215-225.
129. Twigg JP, Irvine DS, Aitken RJ. Oxidative damage to DNA in human spermatozoa does not preclude pronucleus formation at intracytoplasmic sperm injection. *Hum Reprod.* 1998; 13: 1864-1871.
130. Kodama H, Yamaguchi R, Fukuda J, Kasai H, Tanaka T. Increased oxidative deoxyribonucleic acid damage in the spermatozoa of infertile male patients. *Fertil Steril.* 1997; 68: 519-524.
131. Sun JG, Jurisicova A, Casper RF. Detection of deoxyribonucleic acid fragmentation in human sperm: correlation with fertilization in vitro. *Biol Reprod.* 1997; 56: 602-607.
132. Lewis SE, Aitken RJ. DNA damage to spermatozoa has impacts on fertilization and pregnancy. *Cell Tissue Res.* 2005; 322: 33-41.
133. Sawyer DE, Mercer BG, Wiklendt AM, Aitken RJ. Quantitative analysis of gene-specific DNA damage in human spermatozoa. *Mutat Res.* 2003; 529: 21-34.
134. Aitken RJ, Baker MA. Oxidative stress, sperm survival and fertility control. *Mol Cell Endocrinol.* 2006; 250: 66-69.
135. Sakkas D, Mariethoz E, Manicardi G, Bizzaro D, Bianchi PG, Bianchi U. Origin of DNA damage in ejaculated human spermatozoa. *Rev Reprod.* 1999; 4: 31-37.
136. Shen H, Ong C. Detection of oxidative DNA damage in human sperm and its association with sperm function and male infertility. *Free Radic Biol Med.* 2000; 28: 529-536.
137. Evenson DP, Jost LK, Marshall D, Zinaman MJ, Clegg E, Purvis K, et al. Utility of the sperm chromatin structure assay as a diagnostic and prognostic tool in the human fertility clinic. *Hum Reprod.* 1999; 14: 1039-1049.
138. Spano M, Bonde JP, Hjollund HI, Kolstad HA, Cordelli E, Leter G. Sperm chromatin damage impairs human fertility. The Danish First Pregnancy Planner Study Team. *Fertil Steril.* 2000; 73: 43-50.
139. Carrell DT, Liu L, Peterson CM, Jones KP, Hata-saka HH, Erickson L, et al. Sperm DNA fragmentation is increased in couples with unexplained recurrent pregnancy loss. *Arch Androl.* 2003; 49: 49-55.
140. Agarwal A, Allamaneni SS. The effect of sperm DNA damage on assisted reproduction outcomes. *Minerva Ginecol.* 2004; 56: 235-245.
141. Majai G, Petrovski G, Fesus L. Inflammation and the apopto-phagocytic system. *Immunol Lett.* 2006; 104: 94-101.
142. Hikim AP, Wang C, Lue Y, Johnson L, Wang XH, Swerdloff RS. Spontaneous germ cell apoptosis in humans: evidence for ethnic differences in the susceptibility of germ cells to programmed cell death. *J Clin Endocrinol Metab.* 1998; 83: 152-156.
143. Kierszenbaum AL. Apoptosis during spermatogenesis: the thrill of being alive. *Mol Reprod Dev.* 2001; 58: 1-3.
144. Print CG, Loveland KL. Germ cell suicide: new insights into apoptosis during spermatogenesis. *Bioessays.* 2000; 22: 423-430.
145. Jurisicova A, Lopes S, Meriano J, Oppedisano L, Casper RF, Varmuza S. DNA damage in round spermatids of mice with a targeted disruption of the Pp1c-gamma gene and in testicular biopsies of patients with non-obstructive azoospermia. *Mol Hum Reprod.* 1999; 5: 323-330.
146. Lin WW, Lamb DJ, Wheeler TM, Lipshultz LI, Kim ED. In situ end-labeling of human testicular tissue demonstrates increased apoptosis in conditions of abnormal spermatogenesis. *Fertil Steril.* 1997; 68: 1065-1069.
147. Gorczyca W, Traganos F, Jesionowska H, Darzynkiewicz Z. Presence of DNA strand breaks and increased sensitivity of DNA in situ to denaturation in abnormal hu-

- man sperm cells: analogy to apoptosis of somatic cells. *Exp Cell Res.* 1993; 207: 202-205.
146. Irvine DS, Twigg JP, Gordon EL, Fulton N, Milne PA, Aitken RJ. DNA integrity in human spermatozoa: relationships with semen quality. *J Androl.* 2000; 21: 33-44.
149. Wang X, Sharma RK, Sikka SC, Thomas AJ Jr, Falcone T, Agarwal A. Oxidative stress is associated with increased apoptosis leading to spermatozoa DNA damage in patients with male factor infertility. *Fertil Steril.* 2003; 80: 531-535.
150. Agarwal A, Said TM. Oxidative stress, DNA damage and apoptosis in male infertility: a clinical approach. *BJU Int.* 2005; 95: 503-507.
151. Lewis B, Aitken RJ. A redox-regulated tyrosine phosphorylation cascade in rat spermatozoa. *J Androl.* 2001; 22: 611-622.
152. Twigg J, Fulton N, Gomez E, Irvine DS, Aitken RJ. Analysis of the impact of intracellular reactive oxygen species generation on the structural and functional integrity of human spermatozoa: lipid peroxidation, DNA fragmentation and effectiveness of antioxidants. *Hum Reprod.* 1998; 13: 1429-1436.
153. Curi SM, Ariagno JI, Chenlo PH, Mendeluk GR, Pugliese MN, Sardi Segovia LM, et al. Asthenozoospermia: analysis of a large population. *Arch Androl.* 2003; 49: 343-349.
154. Tomlinson MJ, Barratt CL, Cooke ID. Prospective study of leukocytes and leukocyte subpopulations in semen suggests they are not a cause of male infertility. *Fertil Steril.* 1993; 60: 1069-1075.
155. Yanushpolsky EH, Politch JA, Hill JA, Anderson DJ. Is leukocytospermia clinically relevant? *Fertil Steril.* 1996; 66: 822-825.
156. Arata de Bellabarba G, Tortolero I, Villarreal V, Molina CZ, Bellabarba C, Velazquez E. Nonsperm cells in human semen and their relationship with semen parameters. *Arch Androl.* 2000; 45: 131-136.
157. Lemkecher T, Dartigues S, Vaysse J, Kulski O, Barraud-Lange V, Gattegno L, et al. Leucocytospermia, oxidative stress and male fertility: facts and hypotheses. *Gynecol Obstet Fertil.* 2005; 33: 2-10.
158. Vicino M, Loverro G, Simonetti S, Mei L, Selvaggi L. The correlation between idiopathic leukocytospermia, embryo quality and outcome in the FIVET and ICSI procedures. *Minerva Ginecol.* 1999; 51: 413-420.
159. Berger RE, Karp LE, Williamson RA, Koehler J, Moore DE, Holmes KK. The relationship of pyospermia and seminal fluid bacteriology to sperm function as reflected in the sperm penetration assay. *Fertil Steril.* 1982; 37: 557-564.
160. Eggert-Kruse W, Bellmann A, Rohr G, Tilgen W, Runnebaum B. Differentiation of round cells in semen by means of monoclonal antibodies and relationship with male fertility. *Fertil Steril.* 1992; 58: 1046-1055.
161. Menkveld R, Kruger TF. Sperm morphology and male urogenital infections. *Andrologia.* 1998; 30 Suppl 1: 49-53.
162. Thomas J, Fishel SB, Hall JA, Green S, Newton TA, Thornton SJ. Increased polymorphonuclear granulocytes in seminal plasma in relation to sperm morphology. *Hum Reprod.* 1997; 12: 2418-2421.
163. Aziz N, Agarwal A, Lewis-Jones I, Sharma RK, Thomas AJ Jr. Novel associations between specific sperm morphological defects and leukocytospermia. *Fertil Steril.* 2004; 82: 621-627.
164. Fedder J, Askjaer SA, Hjort T. Nonspermatozoal cells in semen: relationship to other semen parameters and fertility status of the couple. *Arch Androl.* 1993; 31: 95-103.
165. Van der Ven HH, Jeyendran RS, Perez-Pelaez M, Al-Hasani S, Diedrich K, Krebs D. Leucospermia and the fertilizing capacity of spermatozoa. *Eur J Obstet Gynecol Reprod Biol.* 1987; 24: 49-52.
166. Lackner JE, Herwig R, Schmidbauer J, Schatzl G, Kratzik C, Marberger M. Correlation of leukocytospermia with clinical infection and the positive effect of anti-inflammatory treatment on semen quality. *Fertil Steril.* 2006; 20: 1-6.
167. Oliva A and Multigner L. Ketotifen improves sperm motility and sperm morphology in male patients with leukocytospermia and unexplained infertility. *Fertile Steril.* 2006; 85: 240-243.
168. Gambera L, Serafini F, Morgante G, Focarelli R, De Leo V, Piomboni P. Sperm quality and pregnancy rate after COX-2 inhibitor therapy of infertile males with abacterial leukocytospermia. *Hum Reprod.* 2007; 22(4): 1047-1051.
169. Pasqualotto FF, Umezu FM, Salvador M, Borges E, Sobreiro BP, Pasqualotto EB. Effect of cigarette smoking on antioxidant levels and presence of leukocytospermia in infertile men: a prospective study. *Fertil Steril.* 2008; 90: 278-283.
170. Young IS, Woodside JV. Antioxidants in health and disease. *J Clin Pathol.* 2001; 54: 176-186.
171. Alvarez JG, Storey BT. Taurine, hypotaurine, epinephrine and albumin inhibit lipid peroxidation in rabbit spermatozoa and protect against loss of motility. *Biol Reprod.* 1983; 29: 548-555.
172. Armstrong JS, Rajasekaran M, Hellstrom WJ, Sikka SC. Antioxidant potential of human serum albumin: role in the recovery of high quality human spermatozoa for assisted reproductive technology. *J Androl.* 1998; 19: 412-419.
173. Sies H. Strategies of antioxidant defense. *Eur J Biochem.* 1993; 215: 213-219.
174. Sikka SC. Oxidative stress and role of antioxidants in normal and abnormal sperm function. *Front Biosci.* 1996; 1: 78-86.
175. Zini A, Fischer MA, Mak V, Phang D, Jarvi K. Catalase-like and superoxide dismutase-like activities in human seminal plasma. *Urol Res.* 2002; 30: 321-323.
176. Nissen HP, Kreysel HW. Superoxide dismutase in human semen. *Klin Wochenschr.* 1983; 61: 63-65.
177. Jeulin C, Soufir JC, Weber P, Laval-Martin D, Calvayrac R. Catalase activity in human spermatozoa and seminal plasma. *Gamete Res.* 1989; 24: 185-196.
178. Alvarez JG, Storey BT. Role of glutathione peroxidase in protecting mammalian spermatozoa from loss of motility caused by spontaneous lipid peroxidation. *Gamete Res.* 1989; 23: 77-90.
179. Li TK. The glutathione and thiol content of mammalian spermatozoa and seminal plasma. *Biol Reprod.* 1975; 12: 641-646.
180. Niki E. Action of ascorbic acid as a scavenger of active and stable oxygen radicals. *Am J Clin Nutr.* 1991; 54: 1119S-1124S.
181. Chow CK. Vitamin E and oxidative stress. *Free*

- Radic Biol Med. 1991; 11: 215-232.
182. Kidd P. Glutathione: Systemic Protectant against Oxidative and Free Radical Damage. *Altern Med Rev*. 1997; 2: 155-177.
183. Gavella M, Lipovac V, Vucic M, Rocic B. Evaluation of ascorbate and urate antioxidant capacity in human semen. *Andrologia*. 1997; 29: 29-35.
184. Ronquist G, Niklasson F. Uridine, xanthine, and urate contents in human seminal plasma. *Arch Androl*. 1984; 13: 63-70.
185. Kodentsova VM, Vrzesinskaya OA, Spirichev VB. Male fertility: a possible role of vitamins. *Ukr Biokhim Zh*. 1994; 66: 17-22.
186. Scandalios JG. Oxidative stress: molecular perception and transduction of signals triggering antioxidant gene defenses. *Braz J Med Biol Res*. 2005; 38: 995-1014.
187. Bauche F, Fouchard MH, Jegou B. Antioxidant system in rat testicular cells. *FEBS Lett*. 1994; 349: 392-396.
188. Yeung CH, Cooper TG, De Geyter M, De Geyter C, Rolf C, Kamischke A, et al. Studies on the origin of redox enzymes in seminal plasma and their relationship with results of in-vitro fertilization. *Mol Hum Reprod*. 1998; 4: 835-839.
189. Ishikawa T, Fujioka H, Ishimura T, Takenake A, Fujisawa M. Increased testicular 8-hydroxy-2'-deoxyguanosine in patients with varicocele. *BJU Int*. 2007; 100: 863-866.
190. Donnelly ET, McClure N, Lewis SE. The effect of ascorbate and alpha-tocopherol supplementation in vitro on DNA integrity and hydrogen peroxide-induced DNA damage in human spermatozoa. *Mutagenesis*. 1999; 14: 505-512.
191. Lewis SE, Sterling ES, Young IS, Thompson W. Comparison of individual antioxidants of sperm and seminal plasma in fertile and infertile men. *Fertil Steril*. 1997; 67: 142-147.
192. Ochsendorf FR, Buhl R, Bastlein A, Beschmann H. Glutathione in spermatozoa and seminal plasma of infertile men. *Hum Reprod*. 1998; 13: 353-359.
193. Said TM, Agarwal A, Sharma RK, Thomas AJ Jr, Sikka SC. Impact of sperm morphology on DNA damage caused by oxidative stress induced by beta-nicotinamide adenine dinucleotide phosphate. *Fertil Steril*. 2005; 83: 95-103.
194. Lissak A, Wiener-Megnazi Z, Reznick AZ, Shnizer S, Ishai D, Grach B, et al. Oxidative stress indices in seminal plasma, as measured by the thermochemiluminescence assay, correlate with sperm parameters. *Fertil Steril*. 2004; 81 Suppl 1: 792-797.
195. Hall JA, Fishel SB, Timson JA, Dowell K, Klentzeris LD. Human sperm morphology evaluation pre- and post-Percoll gradient centrifugation. *Hum Reprod*. 1995; 10: 342-346.
196. Mendiola J, Torres-Cantero AM, Moreno-Grau JM, Ten J, Roca M, Moreno-Grau S, et al. Food intake and its relationship with semen quality: a case-control study. *Fertility Sterility*. 2009; 91(3): 812-818.
197. Mendiola J, Torres-Cantero AM, Vioque J, Moreno-Grau JM, Ten J, Roca M, et al. A low intake of antioxidant nutrients is associated with poor semen quality in patients attending fertility clinics. *Fertility and Sterility*. 2009 (In press).
198. Storey BT, Alvarez JG, Thompson KA. Human sperm glutathione reductase activity in situ reveals limitation in the glutathione antioxidant defense system due to supply of NADPH. *Mol Reprod Dev*. 1998; 49: 400-407.
199. Marklund SL. Human copper-containing superoxide dismutase of high molecular weight. *Proc Natl Acad Sci USA*. 1982; 79: 7634-7638.
200. Marklund SL. Product of extracellular-superoxide dismutase catalysis. *FEBS Lett*. 1985; 184: 237-239.
201. McCord JM, Fridovich I. Superoxide dismutase. An enzymic function for erythrocyte (hemocuprein). *J Biol Chem*. 1969; 244: 6049-6055.
202. Marklund SL, Bjelle A, Elmqvist LG. Superoxide dismutase isoenzymes of the synovial fluid in rheumatoid arthritis and in reactive arthritides. *Ann Rheum Dis*. 1986; 45: 847-851.
203. Marklund SL, Holme E, Hellner L. Superoxide dismutase in extracellular fluids. *Clin Chim Acta*. 1982; 126: 41-51.
204. Oury TD, Day BJ, Crapo JD. Extracellular superoxide dismutase: a regulator of nitric oxide bioavailability. *Lab Invest*. 1996; 75: 617-636.
205. Potts RJ, Notarianni LJ, Jefferies TM. Seminal plasma reduces exogenous oxidative damage to human sperm, determined by the measurement of DNA strand breaks and lipid peroxidation. *Mutat Res*. 2000; 447: 249-256.
206. Lamond S, Watkinson M, Rutherford T, Laing K, Whiting A, Smallwood A, et al. Gene-specific chromatin damage in human spermatozoa can be blocked by antioxidants that target mitochondria. *Reprod Biomed Online*. 2003; 7: 407-418.
207. Nonogaki T, Noda Y, Narimoto K, Umaoka Y, Mori T. Effects of superoxide dismutase on mouse in vitro fertilization and embryo culture system. *J Assist Reprod Genet*. 1992; 9: 274-280.
208. Lasso JL, Noiles EE, Alvarez JG, Storey BT. Mechanism of superoxide dismutase loss from human sperm cells during cryopreservation. *J Androl*. 1994; 15: 255-265.
209. Blum J, Fridovich I. Inactivation of glutathione peroxidase by superoxide radical. *Arch Biochem Biophys*. 1985; 240: 500-508.
210. Baker HW, Brindle J, Irvine DS, Aitken RJ. Protective effect of antioxidants on the impairment of sperm motility by activated polymorphonuclear leukocytes. *Fertil Steril*. 1996; 65: 411-419.
211. Miesel R, Drzejczak PJ, Kurpierz M. Oxidative stress during the interaction of gametes. *Biol Reprod*. 1993; 49: 918-923.
212. Ikeda M, Kodama H, Fukuda J, Shimizu Y, Murata M, Kumagai J, et al. Role of radical oxygen species in rat testicular germ cell apoptosis induced by heat stress. *Biol Reprod*. 1999; 61: 393-399.
213. Kumagai A, Kodama H, Kumagai J, Fukuda J, Kawamura K, Tanikawa H, et al. Xanthine oxidase inhibitors suppress testicular germ cell apoptosis induced by experimental cryptorchidism. *Mol Hum Reprod*. 2002; 8: 118-123.
214. Murawski M, Saczko J, Marcinkowska A, Chwilkowska A, Grybooe M, Banaoe T. Evaluation of superoxide dismutase activity and its impact on semen quality parameters of infertile men. *Folia Histochem Cy-*

- tobiol. 2007; 45, Supp. 1: 123-126.
215. Siciliano L, Tarantino P, Longobardi F, Rago V, De Stefano C, Carpino A. Impaired seminal antioxidant capacity in human semen with hyperviscosity or oligoasthenozoospermia. *J Androl.* 2001; 22: 798-803.
216. Tavilani H, Goodarzi MD, Vaisi-raygani A, Salimi S, Hassanzadeh T. Activity of Antioxidant Enzymes in Seminal Plasma and their Relationship with Lipid Peroxidation of Spermatozoa. *Int Braz J Urol.* 2008; 34 (4): 485-491.
217. Nakamura H, Kimura T, Nakajima A, Shimoya K, Takemura M, Hashimoto K, et al. Detection of oxidative stress in seminal plasma and fractionated sperm from subfertile male patients. *Eur J Obstet Gynecol Reprod Biol.* 2002; 105: 155-160.
218. Lenzi A, Lombardo F, Gandini L, Culasso F, Donde-ro F. Glutathione therapy for male infertility. *Arch Androl.* 1992; 29: 65-68.
219. Parinaud J, Le Lannou D, Vieitez G, Griveau JF, Milhet P, Richoilley G. Enhancement of motility by treating spermatozoa with an antioxidant solution (Sperm-Fit) following ejaculation. *Hum Reprod.* 1997; 12: 2434-2436.
220. Nabil H, Moemen LA and Abu Elela MA. Studying the Levels of Malondialdehyde and Antioxidant Parameters in Normal and Abnormal Human Seminal Plasma. *Aust J Basic Appl Sci.* 2008; 2(3): 773-778.
221. Mostafa T, Tawadrous G, Edward P, Roaia MM, Amer MK, Kader RA, Aziz A. Effect of smoking on seminal plasma ascorbic acid in infertile and fertile males. *Andrologia.* 2006; 38: 221-224.
222. Song GJ, Norku EP, Lewis V. Relationship between seminal ascorbic acid and sperm DNA integrity in infertile men. *Int J Androl.* 2006; 29: 569-575.
223. Oeda T, Henkel R, Ohmori H, Schill WB. Scavenging effect of N-acetyl-L-cysteine against reactive oxygen species in human semen: a possible therapeutic modality for male factor infertility? *Andrologia.* 1997; 29: 125-131.
224. Erkkila K, Hirvonen V, Wuokko E, Parvonen M, Dunkel L. N-acetyl-L-cysteine inhibits apoptosis in human male germ cells in vitro. *J Clin Endocrinol Metab.* 1998; 83: 2523-2531.
225. Irvine DS. Glutathione as a treatment for male infertility. *Rev Reprod.* 1996; 1: 6-12.
226. Muijsers RB, Folkerts G, Henricks PA, Sadeghi-Hashjin G, Nijkamp FP. Peroxynitrite: a two-faced metabolite of nitric oxide. *Life Sci.* 1997; 60: 1833-1845.
227. Wefers H, Sies H. The protection by ascorbate and glutathione against microsomal lipid peroxidation is dependent on vitamin E. *Eur J Biochem.* 1988; 174: 353-357.
228. Giannattasio A, De Rosa M, Smeraglia R, Zarrilli S, Cimmino A, Di Rosario B, et al. Glutathione peroxidase (GPX) activity in seminal plasma of healthy and infertile males. *J Endocrinol Invest.* 2002; 25: 983-986.
229. Garrido N, Meseguer M, Alvarez J, Simón C, Pellicer A, Remoh J. Relationship among standard semen parameters, glutathione peroxidase/glutathione reductase activity, and mRNA expression and reduced glutathione content in ejaculated spermatozoa from fertile and infertile men. *Fertil Steril.* 2004; 82 suppl 3: 1059-1066.
230. Williams AC, Ford WCL. Functional significance of the pentose phosphate pathway and glutathione reductase in the antioxidant defenses of human sperm. *Biol Reprod.* 2004; 71(4): 1309-1316.
231. Aitken J, Fisher H. Reactive oxygen species generation and human spermatozoa: the balance of benefit and risk. *Bioessays.* 1994; 16: 259-267.
232. McCall TB, Boughton-Smith NK, Palmer RM, Whittle BJ, Moncada S. Synthesis of nitric oxide from L-arginine by neutrophils. Release and interaction with superoxide anion. *Biochem J.* 1989; 261: 293-296.
233. Clancy RM, Leszczynska-Piziak J, Abramson SB. Nitric oxide, an endothelial cell relaxation factor, inhibits neutrophil superoxide anion production via a direct action on the NADPH oxidase. *J Clin Invest.* 1992; 90: 1116-1121.
234. Jeyendran RS, Van der Ven HH, Perez-Pelaez M, Crabo BG, Zaneveld LJ. Development of an assay to assess the functional integrity of the human sperm membrane and its relationship to other semen characteristics. *J Reprod Fertil.* 1984; 70: 219-228.
235. Lewis SE, Donnelly ET, Sterling ES, Kennedy MS, Thompson W, Chakravarthy U. Nitric oxide synthase and nitrite production in human spermatozoa: evidence that endogenous nitric oxide is beneficial to sperm motility. *Mol Hum Reprod.* 1996; 2: 873-878.
236. Oettl K, Hallström S, Jürgens G, Fuchs D, Blomster H, Schmut O, et al. Reduced and oxidized forms of human serum albumin in sports, aging and disease. *FEBS Journal.* 2005; 272.
237. Eggert-Kruse W, Zwick EM, Batschulat K, Rohr G, Armbruster FP, Petzoldt D, et al. Are zinc levels in seminal plasma associated with seminal leukocytes and other determinants of semen quality? *Fertil Steril.* 2002; 77: 260-269.
238. Prasad AS. Zinc: an overview. *Nutrition.* 1995; 11: 93-99.
239. Stephenson JL, Brackett BG. Influences of zinc on fertilisation and development of bovine oocytes in vitro. *Zygote.* 1999; 7: 195-201.
240. Gavella M, Lipovac V. In vitro effect of zinc on oxidative changes in human semen. *Andrologia.* 1998; 30: 317-323.
241. Bedwal RS, Bahuguna A. Zinc, copper and selenium in reproduction. *Experientia.* 1994; 50: 626-640.
242. Lewis-Jones DI, Aird IA, Biljan MM, Kingsland CR. Effects of sperm activity on zinc and fructose concentrations in seminal plasma. *Hum Reprod.* 1996; 11: 2465-2467.
243. Chia SE, Ong CN, Chua LH, Ho LM, Tay SK. Comparison of zinc concentrations in blood and seminal plasma and the various sperm parameters between fertile and infertile men. *J Androl.* 2000; 21: 53-57.
244. Hidioglou M, Knipfel JE. Zinc in mammalian sperm: a review. *J Dairy Sci.* 1984; 67: 1147-1156.
245. Prasad AS. Clinical, endocrinological and biochemical effects of zinc deficiency. *Clin Endocrinol Metab.* 1985; 14: 567-589.
246. Davies S. Zinc, nutrition and health. In: Bland J. Editor. *Yearbook of Nutritional Medicine.* New Canaan: Keats Publishing; 1985; 113-152.
247. Abbasi AA, Prasad AS, Rabbani P, DuMouchelle E. Experimental zinc deficiency in man. Effect on testicular function. *J Lab Clin Med.* 1980; 96: 544-550.
248. Piese J. Zinc and Human Male Infertility. *Int Clin Nutr.* 1983; Rev 3: 4-6.

249. Stoltenberg M, Sorensen MB, Danscher G, Juhl S, Andreasen A, Ernst E. Autometallographic demonstration of zinc ions in rat sperm cells. *Mol Hum Reprod*. 1997; 3: 763-767.
250. Fujimori O, Tsukise A, Yamada K. Histochemical demonstration of zinc in rat epididymis using a sulphide-silver method. *Histochemistry*. 1988; 88: 469-473.
251. Stoltenberg M, Ernst E, Andreasen A, Danscher G. Histochemical localization of zinc ions in the epididymis of the rat. *Histochem J*. 1996; 28: 173-185.
252. Ali H, Ahmed M, Baig M, Ali M. Relationship of zinc concentrations in blood and seminal plasma with various semen parameters in infertile subjects. *Pak J Med Sci*. 2007; 23(1): 111-114.
253. Ebisch IMW, Thomas CMG, Peters WHM, Braat DDM, Steegers-Theunissen RPM. The importance of folate, zinc and antioxidants in the pathogenesis and prevention of subfertility. *Hum Reprod Update*. 2007; 13(2): 163-174.
254. Young SS, Eskenazi B, Marchetti FM, Block G, Wyrobek AJ. The association of folate, zinc and antioxidant intake with sperm aneuploidy in healthy non-smoking men. *Hum Reprod*. 2008; 23(5): 1014-1022.
255. Packer L. Vitamin E cycle and benefits of health. In: Yagi K editor. *Active Oxygen, Lipid peroxides and Antioxidants*. Boca Raton FL. CRC Press; 1993; 247-262.
256. Green MH, Lowe JE, Waugh AP, Aldridge KE, Cole J, Arlett CF. Effect of diet and vitamin C on DNA strand breakage in freshly-isolated human white blood cells. *Mutat Res*. 1994; 316: 91-102.
257. Fraga CG, Motchnik PA, Shigenaga MK, Helbock HJ, Jacob RA, Ames BN. Ascorbic acid protects against endogenous oxidative DNA damage in human sperm. *Proc Natl Acad Sci USA*. 1991; 88: 11003-11006.
258. Hughes CM, Lewis SE, McKelvey-Martin VJ, Thompson W. The effects of antioxidant supplementation during Percoll preparation on human sperm DNA integrity. *Hum Reprod*. 1998; 13: 1240-1247.
259. Hughes CM, McKelvey-Martin VJ, Lewis SE. Human sperm DNA integrity assessed by the Comet and ELISA assays. *Mutagenesis*. 1999; 14: 71-75.
260. Brigelius-Flohe R, Traber MG. Vitamin E: function and metabolism. *Faseb J*. 1999; 13: 1145-1155.
261. Burton GW, Joyce A, Ingold KU. Is vitamin E the only lipid-soluble, chain-breaking antioxidant in human blood plasma and erythrocyte membranes? *Arch Biochem Biophys*. 1983; 221: 281-290.
262. Bolle P, Evandri MG, Saso L. The controversial efficacy of vitamin E for human male infertility. *Contraception*. 2002; 65: 313-315.
263. Esterbauer H, Dieber-Rotheneder M, Striegl G, Waeg G. Role of vitamin E in low-density lipoprotein. *Am J Clin Nutr*. 1991; 53: 314S-321S.
264. Ingold KU, Webb AC, Witter D, Burton GW, Metcalfe TA, Muller DP. Vitamin E remains the major lipid-soluble, chain-breaking antioxidant in human plasma even in individuals suffering severe vitamin E deficiency. *Arch Biochem Biophys*. 1987; 259: 224-225.
265. Kamal-Eldin A, Appelqvist LA. The chemistry and antioxidant properties of tocopherols and tocotrienols. *Lipids*. 1996; 31: 671-701.
266. Evans HM, Bishop KS. On the Existence of a Hitherto Unrecognized Dietary Factor Essential for Reproduction. *Science*. 1922; 56: 650-651.
267. Comhaire FH, Christophe AB, Zalata AA, Dhooge WS, Mahmoud AM, Depuydt CE. The effects of combined conventional treatment, oral antioxidants and essential fatty acids on sperm biology in subfertile men. *Prostaglandins Leukot Essent Fatty Acids*. 2000; 63: 159-165.
268. Dawson EB, Harris WA, Teter MC, Powell LC. Effect of ascorbic acid supplementation on the sperm quality of smokers. *Fertil Steril*. 1992; 58: 1034-1039.
269. Yamamoto K, Niki E. Interaction of alpha-tocopherol with iron: antioxidant and prooxidant effects of alpha-tocopherol in the oxidation of lipids in aqueous dispersions in the presence of iron. *Biochim Biophys Acta*. 1988; 958: 19-23.
270. Akmal M, Qadri JQ, Al-Waili NS, Thangal S, Haq A, Saloom KY. Improvement in human semen quality after oral supplementation of vitamin C. *J Med Food*. 2006; 9: 440-442.
271. Johnson FC. The antioxidant vitamins. *CRC Crit Rev Food Sci Nutr*. 1979; 11: 217-309.
272. Williams C, Kronfeld D, Hess T, Waldron J, Saker K, Hoffman R, Harris P. Vitamin E Intake and Oxidative Stress in Endurance Horses. *Equine Nutr Phys Soc Proc*. 2003; 18: 134-135.
273. Putnam ME, Comben N. Vitamin E. *Vet Rec*. 1987; 121: 541-545.
274. Suleiman SA, Ali ME, Zaki ZM, el-Malik EM, Nasr MA. Lipid peroxidation and human sperm motility: protective role of vitamin E. *J Androl*. 1996; 17: 530-537.
275. Kessopoulou E, Powers HJ, Sharma KK, Pearson MJ, Russell JM, Cooke ID, et al. A double-blind randomized placebo cross-over controlled trial using the antioxidant vitamin E to treat reactive oxygen species associated male infertility. *Fertil Steril*. 1995; 64: 825-831.
276. Geva E, Bartoov B, Zabludovsky N, Lessing JB, Lerner-Geva L, Amit A. The effect of antioxidant treatment on human spermatozoa and fertilization rate in an in vitro fertilization program. *Fertil Steril*. 1996; 66: 430-434.
277. Nouri M, Ghasemzadeh A, Farzadi L, Shahnazi V, Novin MG. Vitamins C, E and lipid peroxidation levels in sperm and seminal plasma of asthenoteratozoospermic and normozoospermic men. *IJRM*. 2008; 6(1): 1-5.
278. Becker BF. Towards the physiological function of uric acid. *Free Radic Biol Med*. 1993; 14: 615-631.
279. Persson BE, Sjoman M, Niklasson F, Ronquist G. Uridine, xanthine and urate concentrations in prostatic fluid and seminal plasma of patients with prostatitis. *Eur Urol*. 1991; 19: 253-256.
280. Thiele JJ, Friesleben HJ, Fuchs J, Ochsendorf FR. Ascorbic acid and urate in human seminal plasma: determination and interrelationships with chemiluminescence in washed semen. *Hum Reprod*. 1995; 10: 110-115.
281. Dawson EB, Harris WA, Powell LC. Relationship between ascorbic acid and male fertility. *World Rev Nutr Diet*. 1990; 62: 1-26.
282. Chu FF, Rohan de Silva HA, Esworthy RS, Boteva KK, Walters CE, Roses A, et al. Polymorphism and chromosomal localization of the G1-form of human glutathione peroxidase (GPX2) on 14q24.1 by in situ hybridization. *Genomics*. 1996; 32: 272-276.
283. Cohen H. Human sperm glutathione peroxidase

- isolation and characterisation of a unique selenium enzyme. In: Proceeding of the 78th Annual national meeting of the American society for clinical investigation. Washington. DC. 1986; 34.
284. Zhang LP, Maiorino M, Roveri A, Ursini F. Phospholipid hydroperoxide glutathione peroxidase: specific activity in tissues of rats of different age and comparison with other glutathione peroxidases. *Biochim Biophys Acta*. 1989; 1006:140-143.
285. Behne D, Duk M, Elger W. Selenium content and glutathione peroxidase activity in the testis of the maturing rat. *J Nutr* 1986; 116:1442-1447.
286. Behne D, Weiler H, Kyriakopoulos A. Effects of selenium deficiency on testicular morphology and function in rats. *J Reprod Fertil*. 1996; 106: 291-297.
287. McCoy KE, Weswig PH. Some selenium responses in the rat not related to vitamin E. *J Nutr*. 1969; 98: 383-389.
288. Wu S, Oldfield J, Muth O, Whanger P. Effect of selenium on reproduction. In: Proceeding of the Western Section. American Society Of Animal Science. 1969; 85-89.
289. Maiorino M, Wissing JB, Brigelius-Flohe R, Calabrese F, Roveri A, Steinert P, et al. Testosterone mediates expression of the selenoprotein PHGPx by induction of spermatogenesis and not by direct transcriptional gene activation. *FASEB J*. 1998; 12: 1359-1370.
290. Brown DG, Burk RF. Selenium retention in tissues and sperm of rats fed a Torula yeast diet. *J Nutr*. 1973; 103: 102-108.
291. Calvin HI, Cooper GW, Wallace EW. Evidence that selenium in rat sperm is associated with a cysteine-rich structural proteins of the mitochondrial capsule. *Gamete Res*. 1981; 4: 139-145.
292. Roveri A, Maiorino M, Nisii C, Ursini F. Purification and characterization of phospholipid hydroperoxide glutathione peroxidase from rat testis mitochondrial membranes. *Biochim Biophys Acta*. 1994; 1208: 211-221.
293. Hansen JC, Deguchi Y. Selenium and fertility in animals and man--a review. *Acta Vet Scand*. 1996; 37: 19-30.
294. Ursini F, Heim S, Kiess M, Maiorino M, Roveri A, Wissing J, Flohe L. Dual function of the selenoprotein PHGPx during sperm maturation. *Science*. 1999; 285: 1393-1396.
295. Godeas C, Tramer F, Micali F, Soranzo M, Sandri G, Panfili E. Distribution and possible novel role of phospholipid hydroperoxide glutathione peroxidase in rat epididymal spermatozoa. *Biol Reprod*. 1997; 57: 1502-1508.
296. Maiorino M, Bosello V, Ursini F, Foresta C, Garolla A, Scapin M, et al. Genetic variations of gpx-4 and male infertility in humans. *Biol Reprod*. 2003; 68: 1134-1141.
297. Ursini F, Maiorino M, Brigelius-Flohe R, Aumann KD, Roveri A, Schomburg D, et al. Diversity of glutathione peroxidases. *Methods Enzymol*. 1995; 252: 38-53.
298. Lenzi A, Gandini L, Picardo M, Tramer F, Sandri G, Panfili E. Lipoperoxidation damage of spermatozoa polyunsaturated fatty acids (PUFA): scavenger mechanisms and possible scavenger therapies. *Front Biosci*. 2000; 5: E1-E15.
299. Greco E, Iacobelli M, Rienzi L, Ubaldi F, Ferrero S, Tesarik J. Reduction of the incidence of sperm DNA fragmentation by oral antioxidant treatment. *J Androl*. 2005; 26: 349-353.
300. Calvin HI. Comparative labelling of rat epididymal spermatozoa by intratesticularly administered <sup>65</sup>ZnCl<sub>2</sub> and (<sup>35</sup>S) cysteine. *J Reprod Fertil*. 1981; 61: 65-73.
301. Lenzi A, Picardo M, Gandini L, Lombardo F, Terminali O, Passi S, et al. Glutathione treatment of dyspermia: effect on the lipoperoxidation process. *Hum Reprod*. 1994; 9: 2044-2050.
302. Lenzi A, Culasso F, Gandini L, Lombardo F, Dondero F. Placebo-controlled, double-blind, cross-over trial of glutathione therapy in male infertility. *Hum Reprod*. 1993; 8: 1657-1662.
303. Gil-Villa AM, Cardona-Maya W, Agarawal A, Sharma R, Cadavid A. Role of male factor in early recurrent embryo loss: do antioxidants have any effect?. *Fertil Steril*. 2009; 92(2): 565-571.
304. Piomboni P, Gambera L, Serafini F, Campanella J. After natural anti-oxidant treatment of asthenoteratospermic men with leukocytospermia. *Asian J Androl*. 2008; 10(2): 201-206.
305. Paradiso Galatioto G, Luca Gravina G, Angelozzi G, Sacchetti A, Innominato PF, Pace G, et al. May antioxidant therapy improve sperm parameters of men with persistent oligospermia after retrograde embolization for varicocele?. *World J Urol*. 2008, 26: 97-102.
306. Omu AE, Al-Azemi MK, Kehinde EO, Anim JT, Oriowo MA, Mathew TC. Indications of the Mechanisms Involved in Improved Sperm Parameters by Zinc Therapy. *Med Princ Pract*. 2008; 17: 108-116.
307. Jang HY, Kim SJ, Kim JT, Park CK, Cheong HT, Lee HK, et al. Effects of Antioxidants on Sperm Motility During in Vitro Storage of Boar Semen. *Kor J Gerontol*. 2006; 16(6): 47-51.